arXiv:1003.0743v2 [quant-ph] 21 Jul 2011

A GENERALIZED ANALYTICAL MECHANICS IN WHICH QUANTUM PHENOMENA APPEAR

Masao Yasuda

myasuda@mbn.nifty.com

Analytical mechanics of a spinless massive particle is generalized to that in which quantum phenomena appear. This generalization is accomplished with two keys. The first is the extended action of the diffeomorphism group on the covariant tensor fields of order two, with which the classical Hamilton-Jacobi eq. of the particle is modified. The other is the second-order Lagrangian $L(\mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}})$, with which the Lagrangian formalism consistent with the modified Hamilton-Jacobi eq. is constructed. In the thus obtained formulation, i) the particle picture is held at all times, ii) the discrete energy levels of the system of a particle in a confining potential are the same as those of quantum mechanics, iii) the particle distribution in an ensemble differs slightly from that of quantum mechanics, which however does not mean the incompetence of the present formulation because the precision of the experimental data up to date on the particle distribution is not enough to exclude the present formulation.

Keywords: extensions of the diffeomorphism groups; higher-order analytical mechanics; quantum Hamilton-Jacobi equation; Bohm's interpretation.

PACS numbers: 03.65.Ca

1. Introduction

Despite the great success, serious conceptual difficulties remain unresolved in Quantum Mechanics (QM) (e.g., Ref. 1 chap.17). One of them is that the particle picture cannot be maintained in (the Copenhagen interpretation of) QM; the so-called double-slits problem illustrates this difficulty. There have been attempts to recover the particle picture in QM. Notable is the Bohm's interpretation (B.I.), which allows a realistic particle picture in non-relativistic QM. However, there arise other difficulties in B.I. Most serious would be that it does not work for a relativistic spinless particle (for B.I., see Ref. 2 and references therein).[a]

In Ref. 3, Faraggi and Matone made a breakthrough along the line of B.I. They derived the field eq. used in B.I., which we call the Quantum Stationary Hamilton-Jacobi Equation (QSHJE), modifying the classical Hamilton-Jacobi equation (H-J eq.) with what they call 'the Equivalence Principle', which states that all physical systems can be connected by a coordinate transformation to the free one with vanishing energy. (In classical analytical mechanics (CM), if the energy E of the system vanish: $E = 0$, then $\mathcal{S}_0(x) = const.$, where $\mathcal{S}_0(x)$ is the reduced action. Accordingly, we cannot transform the system to the $E \neq 0$

---

[a] Although talking about the relativistic trajectory goes beyond the scope of the present paper, we can show that the argument against the possibility of introducing the trajectory to a relativistic spinless massive particle in B.I. does not apply to the present formulation; see §4.

system with the time independent coordinate transformation.) To be noted are i) they obtained the QSHJE not from the Schrödinger eq. (SE) of QM (in B.I., it was obtained from SE) but from the H-J eq. of CM, and ii) discrete energy levels appear without recourse to the axiomatic interpretation of the wave function of QM. They appear, in one-dimensional (1-D) space, from the requirement that $e^{2i\mathcal{S}_0(x)/\hbar}$, where $\mathcal{S}_0(x)$ is a solution of the QSHJE, be continuous at $x = \{\pm\infty\}$ in the projective geometrical sense. Their formulation however lacks, as a CM, the Lagrangian formalism in which the trajectory is determined.

In Ref. 4, Bouda determined the 1-D Lagrangian L consistent with 1-D QSHJE in the form: $L(q,\dot{q},\ddot{q},\dddot{q})$ and derived the 1-D Euler-Lagrange eq. (E-L eq.) from it. Bouda determined them considering that the L should depend not only on q, $\dot{q}$ but also on $\ddot{q},\dddot{q},\cdots$ ($\dot{q} = dq/dt$ and so on) because 1-D QSHJE is a third order ordinary differential eq.

Hinted at by their works, we formulate a quantum theory of a non-relativistic spinless massive particle as a generalized, or modified, CM. It does not suffer from the difficulties of B.I. (§4).

We outline our formulation.

Our formulation consists of the H-J formalism and the Lagrangian formalism (the latter is transformable to the Hamiltonian formalism; Appendix C). The H-J formalism is based on the QSHJE:

$$\frac{\hbar^2}{2m}\frac{\nabla^2 R}{R} - \frac{1}{2m}(\partial_i \mathcal{S}_0)^2 - V + E = 0\ , \quad \frac{\partial}{\partial x^i}(\frac{R^2}{m}\partial_i \mathcal{S}_0) = 0\ , \tag{1.1}$$

where $R \in \mathbb{R}$ and $\mathcal{S}_0 \in \mathbb{R}$ are unknown functions. As a partial differential eq., the solution gives a momentum ($\nabla\mathcal{S}_0$) *field*. In the region where $-\hbar^2\nabla^2 R/2mR$, which is called the quantum potential (QP),[2] is not ignorable, the $\nabla\mathcal{S}_0$ can be wave-like if we give suitable initial conditions. The Lagrangian formalism is based on the semiclassical Lagrangian (in principle, we use, as coordinate variables, x and its variants in the H-J formalism, q and its variants in the Lagrangian formalism):

$$L = \frac{1}{2}m\dot{\mathbf{q}}^2 - \frac{\hbar^2(1-\tilde{\boldsymbol{\varepsilon}}(\mathbf{q}))}{2m}\frac{\ddot{\mathbf{q}}^2}{\dot{\mathbf{q}}^4} - V(\mathbf{q})\ , \tag{1.2}$$

where $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q}) << 1$ is determined, in principle, to make $\mathcal{S}_0(\mathbf{x}) - Et = \int^{\mathbf{x},t} L d\tilde{t}$ be true for a solution $\mathcal{S}_0(\mathbf{x})$ of (1.1). Accordingly the $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q})$ makes it possible to determine the trajectory even in the interference region. The $\ddot{\mathbf{q}}^2/\dot{\mathbf{q}}^4$ makes the classical trajectory wavy. The second term of the RHS corresponds to the QP appeared in (1.1). The Lagrangian formalism is effective at classical and semiclassical levels; at the quantum level, our formulation has no Lagrangian (§2.2).

We derived (1.1) modifying the classical H-J eq. with the assumption that the action of the diffeomorphism group on the covariant tensor fields of order two is extended. We note i) although (1.1) is obtained from the SE of QM replacing the wave function Ψ in it with $Re^{i\mathcal{S}_0/\hbar}$ and taking the real and imaginary part of the resulting eq. (e.g., Ref. 2), we do not use this method since, for us, QM is not a starting point but a goal, ii) as was mentioned above, Faraggi and Matone derived (1.1) with ‘the Equivalence Principle’.

We determined our Lagrangian to the form of (1.2) as a quantity which makes $\mathcal{S}_0(\mathbf{x}) - Et = \int^{\mathbf{x},t} L d\tilde{t}$ true for a solution $\mathcal{S}_0$ of (1.1). We note i) the 1-D Lagrangian of Bouda[4] differs from the 1-D form of (1.2); there are some differences between his and our method used to derive the Lagrangian, ii) the Lagrangian formalism is constructed on the higher-order-analytical mechanics, for which see Ref. 5, 6 and references therein.

Quantum phenomena appear as follows.

• Uncertainty relation: In a 1-D square potential well, the momentum $p := \nabla \mathcal{S}_0$ constructed from the solution of the 1-D version of (1.1) undulates. Moreover, the narrower the well, the larger the amplitude of the undulation, which corresponds to the uncertainty relation between position and momentum of QM.

• Energy and angular momentum quantizations: In 1-D space, energy quantization appears from the requirement that $e^{2i\mathcal{S}_0(x)/\hbar}$ is continuous at $x = \{\pm\infty\}$ for a solution $\mathcal{S}_0(x)$ of (1.1), which is equivalent, if the V is the confining one, for the second order linear differential eq. corresponding to (1.1): $-(\hbar^2/2m)(d^2\Psi/dx^2) + (E-V)\Psi = 0$ to have a L^2-solution. Since this eq. is the same as SE, we see that the energy of the system is quantized to the same levels as those of QM.[3] In 3-D space, the energy, the angular momentum too, is quantized by the 3-D version of the 1-D requirement.

• Tunneling: In our formulation, the kinetic energy of the particle depends not only on $\dot{\mathbf{q}}$ but also on $\ddot{\mathbf{q}}$ and $\dddot{\mathbf{q}}$. Accordingly, even if the potential hill is higher than the kinetic energy of the incoming particle, the velocity can be non-zero real: $\dot{\mathbf{q}}^2 > 0$ at the top of the hill.

• Interference: We consider the double-slits experiment for concreteness. In which, the interference pattern on the screen appears as a result of undulating trajectories. The trajectory in the interference region is determined using the formula connecting $\nabla\mathcal{S}_0$ and $(\dot{\mathbf{q}}, \ddot{\mathbf{q}}, \dddot{\mathbf{q}})$: $\nabla\mathcal{S}_0(\mathbf{x}) = m\dot{\mathbf{q}} + (\hbar^2/m)(\dddot{\mathbf{q}}/\dot{\mathbf{q}}^4) + \cdots|_{\mathbf{q}=\mathbf{x}}$ (see (2.36)). We regard this formula as a trajectory determining eq. after inserting the $\nabla\mathcal{S}_0$ constructed from the solution of (1.1) into the LHS. The effect of the double-slits manifests in the functional form of the $\nabla\mathcal{S}_0$. We cannot use the E-L eq. for the present purpose because we cannot give the exact form of $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q})$ in (1.2).

We note above reasoning for 1-D energy quantization is essentially the same as that in Ref. 3. We however repeat it for completeness. Other reasoning would be found nowhere.

Our formulation predicts that the density distribution (dens. distr.) of *massive* particles in an ensemble deviates from $|\Psi|^2$ (by a few percent in the experimental setting of §3.3), which however does not contradict existing data supporting QM because the precision of the data is not enough to exclude our prediction. For example, the experimental data of the dens. distr. of electrons Fresnel-diffracted by the electron biprism, [7] [8] [9] which is supposed to support QM's prediction, does not exclude our formulation because the experimental error (here we mean the difference between the measured and quantum mechanically calculated dens. distr.) by more than several percent [9] is larger than the expected difference between QM's and our prediction by a few percent. Thus, we see that our formulation has a chance to be verified by experiments. (Casual inspection makes us feel that the data support our formulation; see the last paragraph of §3.3.)

The organization of the present paper is as follows. In §2.1, we derive the QSHJE. In §2.2, we construct the Lagrangian formalism. In §3, we show that above mentioned quantum phenomena appear. In particular, in §3.3, we show that the dens. distr. of diffracted electrons differs from that of QM. In §4, we describe how our formulation differs from B.I. In appendix, we put background materials and lengthy formula derivations.

2. Formalism

2.1. *Quantum Hamilton-Jacobi equation*

We first modify the relativistic H-J eq. assuming the extended diffeomorphism action. Then, we take the non-relativistic limit of the modified H-J eq.

The extended action of the diffeomorphism group of 1-D projective space $\mathbb{RP}^1$: $\mathrm{Diff}(\mathbb{RP}^1)$ upon the covariant tensor fields of order two on $\mathbb{RP}^1$: $\mathcal{F}_2(\mathbb{RP}^1)$ is possible; see Ref. 10 and Appendix A. The (not-extended) action T of $\mathrm{Diff}(\mathbb{RP}^1)$ on $\mathcal{F}_2(\mathbb{RP}^1)$ is given as $T : \mathrm{Diff}(\mathbb{RP}^1) \times \mathcal{F}_2(\mathbb{RP}^1) \to \mathcal{F}_2(\mathbb{RP}^1),\ (y^{-1}, \phi(y)) \mapsto T_{y^{-1}}\phi(y) = \phi(y(x))(dy/dx)^2$. While the extended action (we write it as $T_{(dy^{-1},\, y^{-1})}$) is given as

$$T_{(dy^{-1},\, y^{-1})}\phi(y) = (\frac{dy}{dx})^2\phi(y(x)) + \alpha\{y(x), x\} \ , \tag{2.1}$$

where $\alpha = const. \in \mathbb{R}$ and $\{y(x), x\} = \frac{y'''}{y'} - \frac{3}{2}(\frac{y''}{y'})^2$ is the Schwarzian derivative ($y' = \partial y/\partial x$ and so on). We call this action the extended action of the diffeomorphism group or the action of the extended diffeomorphism group.

The extended action of the diffeomorphism group of 4-D Minkowski spacetime $\mathcal{M}^4$ on the covariant tensor fields of order two on $\mathcal{M}^4$: $\mathsf{T}_{\alpha\beta}(\mathcal{M}^4)$ is also possible (the $\mathcal{F}_2(\mathbb{RP}^1)$ is identified as a restriction of $\mathsf{T}_{\alpha\beta}(\mathcal{M}^4)$ to 1-D space). Indeed, we constructed the extended action 'on' the H-J eq.; see Appendix B. We consider the diffeomorphism of $\mathcal{M}^4$ as the one between patches not including infinity. We therefore do not care about the global topology of the base manifold. With the extended action, the H-J eq. (our Minkowski metric is $\mathrm{diag}(1,-1,-1,-1)$):

$$(\frac{\partial \mathcal{S}}{\partial \mathbf{x}} - \frac{e}{c}\mathbf{A})^2 - \frac{1}{c^2}(\frac{\partial \mathcal{S}}{\partial t} + eA^0)^2 + m^2c^2 = 0 \ , \tag{2.2}$$

where $\mathcal{S}$ is the relativistic action, e is the electric charge of the particle, m is the mass of the particle, c is the velocity of light, and $(A^0, \mathbf{A})$ is the electromagnetic potential, is modified to what we call the relativistic quantum H-J eq.:

$$(\partial_\mu \mathcal{S} + \frac{e}{c}A_\mu)(\partial^\mu \mathcal{S} + \frac{e}{c}A^\mu) = m^2(c^2 + Q), \quad Q = \frac{\hbar^2 \partial_\mu \partial^\mu R}{Rm^2} \tag{2.3a}$$

$$\partial_\mu j^\mu = 0, \qquad j^\mu = -\frac{R^2}{m}(\partial^\mu \mathcal{S} + \frac{e}{c}A^\mu) \ , \tag{2.3b}$$

where μ runs over $0 \sim 3$, and R is a function determined solving the equations.

Taking non-relativistic limit, that is, setting $\mathcal{S} = \mathcal{S}_{nr} - mc^2 t$ ($\mathcal{S}_{nr}$ is the non-relativistic action) and ignoring $\bullet/c^2$ terms (we consider only the case: $A^1 = A^2 = A^3 = 0$ and A^0 is time-independent; in this case, R is time-independent and $\mathcal{S}_{nr}$ is decomposed as $\mathcal{S}_{nr}(\mathbf{x}, t) = \mathcal{S}_0(\mathbf{x}) - Et$, where $\mathcal{S}_0$ is the reduced action), we have the non-relativistic quantum H-J eq.

$$\frac{\hbar^2}{2m}\frac{\nabla^2 R}{R} - \frac{1}{2m}(\partial_i \mathcal{S}_0)^2 - eA^0 + E = 0 \ , \quad \frac{\partial}{\partial x^i}(\frac{R^2}{m}\partial_i \mathcal{S}_0) = 0 \ , \tag{2.4}$$

where i runs over $1, 2, 3$. If space dimension is one, (2.4) is simplified, using the formula in footnote x, to

$$\frac{1}{2m}(\frac{\partial \mathcal{S}_0}{\partial x})^2 + \frac{\hbar^2}{4m}\{\mathcal{S}_0, x\} + V(x) - E = 0 \ , \tag{2.5}$$

where $V(x) = eA^0(x)$. We call (2.4) and (2.5) 3-D QSHJE and 1-D QSHJE respectively after Ref. 3.[b] [c] We call $\hbar^2\nabla^2 R/(2mR)$ and $\hbar^2\{\mathcal{S}_0, x\}/(4m)$ 3-D and 1-D quantum potential (QP) respectively. Replacing $q^a \to x,\ q^b \to \mathcal{S}_0,\ q^c \to e^{2i\mathcal{S}_0/\hbar}$ in the cocycle condition: $\{q^c, q^a\} = (\partial q^b/\partial q^a)^2\{q^c, q^b\} + \{q^b, q^a\}$, which is equivalent to $d(g^{-1}h^{-1}) = dg^{-1} + T_{g^{-1}}dh^{-1}$ on p.30 (these replacements are allowed because $x \mapsto \mathcal{S}_0(x)$, $\mathcal{S}_0 \mapsto e^{2i\mathcal{S}_0/\hbar}$, etc. are at least local diffeomorphisms; see footnote v), we rewrite (2.5) as

$$V(x) - E = -\frac{\hbar^2}{4m}\{e^{\frac{2i}{\hbar}\mathcal{S}_0}, x\}\ . \tag{2.6}$$

Using the theorem: 'Let the two linearly independent solutions of $d^2z/dx^2 + u(x)z = 0$ be z_1, z_2. Then, $\{z_2/z_1, x\} = \{(Az_1 + Bz_2)/(Cz_1 + Dz_2), x\} = 2u(x)$, where $A, \cdots, D = const. \in \mathbb{C}$ and $AD - BC \neq 0$.' (e.g., Ref. 10), we solve the 1-D QSHJE (2.6) as

$$e^{\frac{2i}{\hbar}\mathcal{S}_0} = \frac{A\Psi^D + B\Psi}{C\Psi^D + D\Psi}\ , \tag{2.7}$$

where Ψ and Ψ^D are two linearly independent solutions of the corresponding (having the same $V(x)$ and E) second-order linear differential eq.: $d^2z/dx^2 - (2m/\hbar^2)(V(x) - E)z = 0$. Since this eq. is equivalent to the SE of QM, we henceforth call this eq. SE. Keep in mind however that in our formulation SE is used only as a mathematical tool to solve (2.5) (or its 3-D version (2.4) as is seen on p.7). We do not give any physical meaning such as '$|\Psi(x)|^2$ represents the probability density of finding a particle at x ' to Ψ or Ψ^D. Physics is discussed with $\mathcal{S}_0$.

The momentum $p := \partial\mathcal{S}_0/\partial x$ is obtained differentiating both sides of (2.7) with respect to q as

$$p = \frac{i}{2}\frac{\hbar(AD - BC)W}{(A\Psi^D + B\Psi)(C\Psi^D + D\Psi)} = \frac{\frac{i\hbar(AD-BC)W}{2AC}}{(\Psi^D + \frac{AD+BC}{2AC}\Psi)^2 - (\frac{AD-BC}{2AC})^2\Psi^2}, \tag{2.8}$$

where W is the Wronskian: $W = \Psi'\Psi^D - \Psi^{D\prime}\Psi = const. \neq 0$. We require p to be real because p is a classically observable quantity. We make (2.8) a little readable. We set, without losing generality, $\Psi, \Psi^D \in \mathbb{R}$ and $AD - BC = \pm 2i$ for convenience. Then, with $l_1 = i(AD - BC)/2AC = \mp 1/AC \in \mathbb{R}\backslash\{0\}$ and $l_2 = (AD + BC)/2AC \in \mathbb{R}$, (2.8) is transformed as follows with p being kept real:

$$p = \frac{\partial\mathcal{S}_0}{\partial x} = \frac{\hbar W l_1}{(\Psi^D + l_2\Psi)^2 + l_1^2\Psi^2}\ . \tag{2.9}$$

In accordance with the appearance in (2.6) of the Schwarzian derivative, which lives in 1-D projective space $\mathbb{RP}^1$, we put a requirement on the $\mathcal{S}_0$ that, as a physically acceptable solution,

$$\text{the } e^{2i\mathcal{S}_0/\hbar} \text{ is continuous on } \hat{\mathbb{R}} := \mathbb{R} \cup \{\pm\infty\} (\approx \mathbb{RP}^1). \tag{2.10}$$

---

[b] As was mentioned in §1, Faraggi and Matone derived (2.5) in Ref. 3 through a different path from ours, through what they call 'the Equivalence Principle'.

[c] In writing this and following paragraphs to the end of this subsection, corresponding parts of Ref. 3 were helpful.

Since $e^{2i\mathcal{S}_0/\hbar}$ is always continuous (even differentiable) on $\mathbb{R}$ as is seen from (2.9), this requirement is essentially the continuity [d] [e] requirement at $x=\{\pm\infty\}$. Since the $e^{2i\mathcal{S}_0/\hbar}$ is continuous at $x=\{\pm\infty\}$ if and only if $\lim_{x\to+\infty}\mathcal{S}_0(x)-\lim_{x\to-\infty}\mathcal{S}_0(x)=nh/2$ ($n=\pm1,\pm2,\cdots$), (2.10) is equivalent to the requirement:

$$\int d\mathcal{S}_0=\int_{-\infty}^{+\infty}p\,dx=\frac{nh}{2}\ , \tag{2.11}$$

which corresponds to the Bohr-Sommerfeld quantization condition: $\oint p\,dx=\oint d\mathcal{S}_0=nh$. We consider that the $e^{2i\mathcal{S}_0/\hbar}$ is continuous at $x=\{\pm\infty\}$ for $n=\pm\infty$. If we use (2.11) instead of (2.10), we need not refer to the neighborhood of infinity; if the physical system under consideration is of atomic scale, integration interval of (2.11) is taken to be much less than, say, $1\mu m$. Thus, although our formulation is constructed on $\mathbb{RP}^1$, the formulation is indistinguishable from the one on $\mathbb{R}$ on which CM and QM are considered to reside.

In the case of a confining potential, (2.10) is equivalent also to the requirement:

$$\text{The corresponding SE has a } L^2\text{-solution,} \tag{2.12}$$

which is seen as follows. Inserting (2.9) into (2.11) and using a formula: $\int[(\,f'(x)g(x)-f(x)g'(x)\,)/(\,f^2(x)+g^2(x)\,)]dx=\arctan[f(x)/g(x)]$, we rewrite (2.11) as

$$\int_{-\infty}^{+\infty}p\,dx=\int_{-\infty}^{+\infty}\frac{\hbar W l_1}{(\Psi^D+l_2\Psi)^2+l_1^2\Psi^2}dx=-\hbar\arctan\frac{\Psi^D+l_2\Psi}{l_1\Psi}\Big|_{-\infty}^{+\infty}=\frac{nh}{2}\ , \tag{2.13}$$

which is equivalent to require for the Ψ^D/Ψ to be continuous at $x=\{\pm\infty\}$. In the case of a confining potential, the Ψ^D/Ψ is continuous at $x=\{\pm\infty\}$ if and only if corresponding SE has a L^2-solution, which is seen from i) $\sim$ iii) below. Let Ψ_c and Ψ_c^D be two linearly independent solutions of the 1-D SE having a confining potential. Then,

i) The Ψ_c can be an L^2-solution only for discrete values of E. For such E's, the other solution Ψ_c^D diverges for $x=\pm\infty$.[14]

ii) If the Ψ_c is a L^2-solution, Ψ_c^D/Ψ_c is continuous at $x=\{\pm\infty\}$, which is seen from

$$\Psi^D(x)=c\Psi(x)\int_{x_0}^{x}\Psi^{-2}(q)dq+d\Psi(x)\ , \tag{2.14}$$

where $c,d\in\mathbb{C}$ ($c\neq0$) are consts. (the formula (2.14) is obtained solving $W=\Psi'\Psi^D-\Psi^{D\prime}\Psi=const.$ regarded as a differential eq. for Ψ^D with the variation of constants method).

iii) If none of Ψ_c and Ψ_c^D is an L^2-solution, Ψ_c^D/Ψ_c cannot be continuous at $x=\{\pm\infty\}$ because Ψ_c and Ψ_c^D behave for $x\to\pm\infty$ as follows (Ref. 14 chap.3).

---

[d] A function $f(x)\in\hat{\mathbb{R}}$ on $\hat{\mathbb{R}}$ is said to be continuous at $x_0\in\hat{\mathbb{R}}$ if

$$\lim_{x\uparrow x_0}f(x)=\begin{cases}\lim_{x\downarrow x_0}f(x)=f(x_0) & \text{for } f(x_0)\in\mathbb{R}\\ -\lim_{x\downarrow x_0}f(x) & \text{for } \lim_{x\to x_0}f(x)=+\infty \text{ or } -\infty\ .\end{cases}$$

[e] We cannot require differentiability at $x=\{\pm\infty\}$ as is seen from the following example. Consider the particle moving in a non-confining potential. Let the energy of the system be E and the potential approaches to a constant value $V_-<E$ for $x\to-\infty$ and $V_+(\neq V_-)<E$ for $x\to+\infty$. Then, $\lim_{x\to-\infty}\partial\mathcal{S}_0/\partial x=\sqrt{2m(E-V_-)}\neq\sqrt{2m(E-V_+)}=\lim_{x\to+\infty}\partial\mathcal{S}_0/\partial x$.

For $x \to -\infty$, $\Psi_c \to \pm\infty$ and $\Psi_c^D \to 0$; for $x \to +\infty$, $\Psi_c \to 0$ and $\Psi_c^D \to \pm\infty$. Or,
for $x \to -\infty$, $\Psi_c \to 0$ and $\Psi_c^D \to \pm\infty$; for $x \to +\infty$, $\Psi_c \to \pm\infty$ and $\Psi_c^D \to 0$.
Thus, in the case of a confining potential, (2.12) is equivalent to (2.11) or (2.10).

If space dimension is three, the solution set $(R, \mathcal{S}_0)$, where $R, \mathcal{S}_0 \in \mathbb{R}$, of 3-D QSHJE (2.4) is constructed from the solution ψ of the corresponding SE as $\psi = Re^{i\mathcal{S}_0/\hbar}$ since (2.4) is equivalent to the SE with $\psi = Re^{i\mathcal{S}_0/\hbar}$ (e.g., Ref. 2). The momentum $\mathbf{p} := \nabla \mathcal{S}_0$ is given as

$$\mathbf{p} = \nabla \mathcal{S}_0 = \frac{\hbar}{i} \frac{\bar{\psi}\nabla\psi - \psi\nabla\bar{\psi}}{2\psi\bar{\psi}} \ , \tag{2.15}$$

where $\bar{\psi}$ is the complex conjugate of ψ. Of course, (2.15) is valid also in 1-D space. In 1-D space, the p obtained from (2.15) setting $\psi = c_1\Psi + c_2\Psi^D$ is equal to the p obtained from (2.8) with $A = c_1, B = c_2, C = \bar{c_1}$ and $D = \bar{c_2}$. The requirement imposed on 3-D $\mathcal{S}_0$ is given in §3.2.

Example We consider a 1-D free particle of energy $E \geq 0$.

If $E > 0$, we take $\Psi = \cos kx$ and $\Psi^D = \sin kx$, where $k = \sqrt{2mE}/\hbar$, as two linearly independent solutions of the corresponding SE. The momentum p is given, from (2.9), as

$$p = \frac{-\hbar k l_1}{(\sin kx + l_2 \cos kx)^2 + l_1^2 \cos^2 kx} \ . \tag{2.16}$$

If $l_1 = \mp 1$ and $l_2 = 0$, we have the classical value: $p = \pm\sqrt{2mE}$.

If $E = 0$, we take $\Psi = \mathbf{1}$ and $\Psi^D = x$, where $\mathbf{1}$ is a constant having the same dimension as x. The momentum p is given as

$$p := \frac{\partial \mathcal{S}_0}{\partial x} = \frac{-\hbar l_1 \mathbf{1}}{(x + l_2\mathbf{1})^2 + (l_1\mathbf{1})^2} \ . \tag{2.17}$$

The $e^{2i\mathcal{S}_0/\hbar}$ is continuous at $x = \{\pm\infty\}$ since we have $\lim_{x\to+\infty} \mathcal{S}_0(x) - \lim_{x\to-\infty} \mathcal{S}_0(x) = \pm h/2$ from (2.17). The $\mathcal{S}_0$ constructed from $e^{2i\mathcal{S}_0/\hbar} = (Ax + B\mathbf{1})/(Cx + D\mathbf{1})$ is therefore physically acceptable with $A \sim D$ which make p real.

2.2. *Lagrangian formalism*

From the solution $\mathcal{S}_0(\mathbf{x})$ of the QSHJE, we learn the value of the particle's momentum $\mathbf{p} := \nabla\mathcal{S}_0$ at $\mathbf{x}$. From the QSHJE, however, we cannot see whether the particle comes to $\mathbf{x}$ or not, nor see we how $\mathbf{p}$ is related to ($\dot{\mathbf{q}}$, $\ddot{\mathbf{q}}, \cdots$) of the particle ($\nabla\mathcal{S}_0(\mathbf{x}) = m\dot{\mathbf{q}}|_{\mathbf{q}=\mathbf{x}}$ is no longer valid). To see these, we construct the Lagrangian formalism.

We consider, based on a philosophy of CM, the eq. of motion exists as an ordinary differential eq. (ODE) even at the quantum level. Although not all ODEs are obtained with the variational method from the Lagrangian [15], we search for the eq. assuming that our system has the Lagrangian. If not, we will see contradictions somewhere in the eq. searching process. Indeed, we will find that the Lagrangian does not exist beyond the semiclassical region, at least within our theoretical framework.

Our semiclassical Lagrangian cannot be a function of $\mathbf{q}$ and $\dot{\mathbf{q}}$ as in CM. We show the reason. In the Lagrangian formalism, the $\mathbf{p}$ is a function of the trajectory $\mathbf{q}(t)$. That is, the $\mathbf{p}$ at the particle position $\mathbf{q}(t)$ is given as $\mathbf{p}|_{\mathbf{q}(t)} = f(\mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}}, \cdots)|_t \cdots (*1)$. Let us consider a 1-D free particle. Assume (to get a contradiction) that our Lagrangian of which is a function of q and $\dot{q}$: $L(q, \dot{q})$. Then, the E-L eq. is of second order. The trajectory is therefore fixed if two integration parameters, say, particle position q_0 at time t_0: $q_0(t_0)$ and the energy E, are given. The momentum p at any point on the trajectory is also fixed from $(*1)$. However, the formula (2.16) tells us that, even if q_0 and E are given, the p at, say, q_0 is not fixed, which implies that the trajectory is not fixed with $q_0(t_0)$ and E only.

If the E-L eq. is of fourth order, the above problem is resolved because four parameters (x, E, l_1, l_2) necessary to fix the p of (2.16) correspond to four integration parameters of the E-L eq. Thus, we introduce the higher-order Lagrangian $L(q, \dot{q}, \ddot{q}, \cdots)$, from which fourth order E-L eq. is derived. Or we assume that the system described by the QSHJE has the Lagrangian with $\mathcal{S}_0(\mathbf{x}) - Et = \int^{\mathbf{x},t} L(\mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}}, \cdots) d\tilde{t} \cdots (*2)$, where $\mathcal{S}_0(\mathbf{x})$ is a solution of the QSHJE.

We note on a seeming inconsistency in the $(*2)$. In the second (similar for higher) order Lagrangian formalism, the reduced action $\tilde{\mathcal{S}}_0(\mathbf{x}, \dot{\mathbf{x}})$ is obtained as a function of $\mathbf{x}$ and $\dot{\mathbf{x}}$ from $\tilde{\mathcal{S}}_0(\mathbf{x}, \dot{\mathbf{x}}) - Et = \int^{\mathbf{x},\dot{\mathbf{x}},t} L(\mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}}) d\tilde{t} \cdots (*3)$, where $\dot{\mathbf{x}}$ is the velocity at the end point $\mathbf{x}$ (see (C.1), see Ref. 22 §9). Whereas the reduced action $\mathcal{S}_0(\mathbf{x})$ obtained from the QSHJE is a function of $\mathbf{x}$. This seeming inconsistency is understood as follows. In $(*2)$, the integration is carried out along the trajectory. Since the trajectory, which is determined by the E-L eq., depends on initial (ini.) conditions, the $\mathcal{S}_0(\mathbf{x})$ too depends on the ini. conditions of the E-L eq.—we may write $\mathcal{S}_0(\mathbf{x})$ as $\mathcal{S}_0(\mathbf{x}; ic_1, ic_2, \cdots)$, where $ic_1, \cdots$ are ini. conditions. If we take $\mathbf{x}(t)$ and $\dot{\mathbf{x}}(t)$ as parts of the ini. conditions, the $(*2)$ is written in the form of $(*3)$. Thus, we see that $(*2)$ and $(*3)$ are essentially the same; In $(*3)$, parts of the ini. conditions $\mathbf{x}(t)$ and $\dot{\mathbf{x}}(t)$ are explicit, whereas in $(*2)$, ini. conditions other than the $\mathbf{x}(t)$ are implicit.

In the following, we determine the semiclassical Lagrangian of our formulation making use of the higher-order CM (Appendix C). First, we determine the Lagrangian of a free particle in 1-D space. Then, we generalize it to that of a particle in a 3-D potential field.

We expand the would-be 1-D free particle Lagrangian $L = L(q, \dot{q}, \ddot{q}, \dot{\ddot{q}}, \cdots)$ in power series of $\hbar$:

$$L = \frac{1}{2}m\dot{q}^2 + \hbar f_1(\dot{q},\ddot{q},\dddot{q},\ddddot{q},\cdots) + \hbar^2 f_2(\dot{q},\ddot{q},\dddot{q},\ddddot{q},\cdots) + \cdots \ .$$

We assume that the physical parameter (the parameter having dimensions) appearing in each term $f_1, f_2, \cdots$ is only mass m. Then, from the dimensional analysis, we see that f_k's are restricted to

$$f_1 = \frac{c_1\ddot{q}}{\dot{q}} \ , \quad f_2 = \frac{c_{2a}}{m}\frac{\dddot{q}}{\dot{q}^3} \text{ or } f_2 = \frac{c_{2b}}{m}\frac{\ddot{q}^2}{\dot{q}^4} \ , \quad f_k = \frac{c_k}{m^{k-1}}\frac{\ddot{q}^k}{\dot{q}^{3k-2}} \quad (k = 3,4,\cdots) \ , \tag{2.18}$$

where $c_1, c_{2a}, \cdots$ are dimensionless parameters independent from $q, \cdots, \dddot{q}$. Thus, we have

$$L = \frac{1}{2}m\dot{q}^2 + c_1\hbar\frac{\ddot{q}}{\dot{q}} + \frac{c_{2a}\hbar^2}{m}\frac{\dddot{q}}{\dot{q}^3} + \frac{c_{2b}\hbar^2}{m}\frac{\ddot{q}^2}{\dot{q}^4} + \sum_{k=3}^{\infty}\frac{c_k\hbar^k}{m^{k-1}}\frac{\ddot{q}^k}{\dot{q}^{3k-2}} \ . \tag{2.19}$$

The second and higher terms of the RHS as a whole is the Lagrangian side counterpart of the QP appeared in the QSHJE (2.5). We call it too the QP. The $c_1, c_{2a}, \cdots$ are determined by l_1, l_2 and $k = \sqrt{2mE}/\hbar$ of (2.16). Although $c_1, c_3, c_5, \cdots$-terms are not invariant under time reversal, we keep them intact for the moment. The (2.19) is not defined for $\dot{q} = 0$ due to the $\dot{q}$ in denominators, which however causes no problem because the $\dot{q}$ of a solution of the E-L eq. never takes value zero unless $\ddot{q} \equiv 0$ (Appendix E).

The E-L eq., J-O momenta $p_{(1)}$, $p_{(2)}$, $p_{(3)}$, the energy E obtained from (2.19) are (Appendix C and Appendix D)

$$\begin{aligned} 0 = m\ddot{q} &+ \frac{2(3c_{2a}+c_{2b})\hbar^2}{m}\Big(-\frac{\ddddot{q}}{\dot{q}^4} + \frac{8\ddot{q}\dddot{q}}{\dot{q}^5} - \frac{10\ddot{q}^3}{\dot{q}^6}\Big) + \sum_{k=3}^{\infty}\frac{(k-1)c_k\hbar^k}{m^{k-1}} \\ &\times \Big(-\frac{(k-2)k\ddot{q}^{k-3}\dddot{q}^2 + k\ddot{q}^{k-2}\ddddot{q}}{\dot{q}^{3k-2}} + \frac{2(3k-2)k\ddot{q}^{k-1}\dddot{q}}{\dot{q}^{3k-1}} - \frac{(3k-2)(3k-1)\ddot{q}^{k+1}}{\dot{q}^{3k}}\Big) \ . \end{aligned} \tag{2.20}$$

$$\begin{aligned} p_{(1)} = \frac{\partial L}{\partial \dot{q}} - \frac{d}{dt}\frac{\partial L}{\partial \ddot{q}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \dddot{q}} &= m\dot{q} - \frac{2(3c_{2a}+c_{2b})\hbar^2}{m}\frac{\dddot{q}}{\dot{q}^4} + \frac{4(3c_{2a}+c_{2b})\hbar^2}{m}\frac{\ddot{q}^2}{\dot{q}^5} \\ &+ \sum_{k=3}^{\infty}\frac{c_k\hbar^k}{m^{k-1}}\Big(-\frac{k(k-1)\ddot{q}^{k-2}\dddot{q}}{\dot{q}^{3k-2}} + \frac{(3k-2)(k-1)\ddot{q}^k}{\dot{q}^{3k-1}}\Big) \ , \end{aligned} \tag{2.21a}$$

$$p_{(2)} = \frac{\partial L}{\partial \ddot{q}} - \frac{d}{dt}\frac{\partial L}{\partial \dddot{q}} = \frac{c_1\hbar}{\dot{q}} + \frac{(3c_{2a}+2c_{2b})\hbar^2}{m}\frac{\ddot{q}}{\dot{q}^4} + \sum_{k=3}^{\infty}\frac{kc_k\hbar^k}{m^{k-1}}\frac{\ddot{q}^{k-1}}{\dot{q}^{3k-2}} \ , \tag{2.21b}$$

$$p_{(3)} = \frac{\partial L}{\partial \dddot{q}} = \frac{c_{2a}\hbar^2}{m\dot{q}^3} \ . \tag{2.21c}$$

$$\begin{aligned} E = p_{(1)}\dot{q} + p_{(2)}\ddot{q} + p_{(3)}\dddot{q} - L &= \frac{1}{2}m\dot{q}^2 - \frac{2(3c_{2a}+c_{2b})\hbar^2}{m}\frac{\dddot{q}}{\dot{q}^3} + \frac{5(3c_{2a}+c_{2b})\hbar^2}{m}\frac{\ddot{q}^2}{\dot{q}^4} \\ &+ \sum_{k=3}^{\infty}\frac{c_k\hbar^k}{m^{k-1}}\Big(-\frac{k(k-1)\ddot{q}^{k-2}\dddot{q}}{\dot{q}^{3k-3}} + \frac{(3k-1)(k-1)\ddot{q}^k}{\dot{q}^{3k-2}}\Big) \ . \end{aligned} \tag{2.22}$$

Note i) the E-L eq. is fixed to the fourth order only from the dimensional analysis, ii) the c_1 does not appear in the E-L eq., $p_{(1)}$ and E.

The momentum $p_{EL} := \partial \mathcal{S}_0(x)/\partial x$ obtained from (2.19) is

$$\begin{aligned} p_{EL} &= p_{(1)} + p_{(2)}\frac{\ddot{q}}{\dot{q}} + p_{(3)}\frac{\dddot{q}}{\dot{q}} \\ &= p_{(1)} + c_1\hbar\frac{\ddot{q}}{\dot{q}^2} + \frac{c_{2a}\hbar^2}{m}\frac{\dddot{q}}{\dot{q}^4} + \frac{(3c_{2a}+2c_{2b})\hbar^2}{m}\frac{\ddot{q}^2}{\dot{q}^5} + \sum_{k=3}^{\infty}\frac{kc_k\hbar^k}{m^{k-1}}\frac{\ddot{q}^k}{\dot{q}^{3k-1}} \ , \end{aligned} \tag{2.23}$$

which is obtained as follows. We know (Appendix C):

$$\mathcal{S} = \int L(q,\dot{q},\ddot{q},\dddot{q})dt \ , \quad E = p_{(1)}\dot{q} + p_{(2)}\ddot{q} + p_{(3)}\dddot{q} - L \ . \tag{2.24}$$

From (2.24), along the solution path, we have

$$\begin{aligned} d\mathcal{S} &= Ldt = (p_{(1)}\dot{q} + p_{(2)}\ddot{q} + p_{(3)}\dddot{q} - E)dt \\ &= p_{(1)}dq + p_{(2)}d\dot{q} + p_{(3)}d\ddot{q} - Edt = (p_{(1)} + p_{(2)}\frac{\ddot{q}}{\dot{q}} + p_{(3)}\frac{\dddot{q}}{\dot{q}})dq - Edt \ . \end{aligned} \tag{2.25}$$

Comparing (2.25) with $d\mathcal{S} = d\mathcal{S}_0 - Edt = pdq - Edt$ of CM, we see that $p_{EL} = p_{(1)} + p_{(2)}\ddot{q}/\dot{q} + p_{(3)}\dddot{q}/\dot{q}$. Thus, we have (2.23) (the $p_{(1)}$, which is const. from (C.6), is left intact for later convenience).

For the E-L eq. (2.20) to be the eq. of motion, it has to be $3c_{2a} + c_{2b} < 0$. Otherwise, a wavy linear trajectory required for the p_{EL} to correspond to the p of (2.16) cannot be a solution of (2.20). We call p of (2.16) p_{QHJ} henceforth.

If $c_2 := 3c_{2a} + c_{2b} < 0$, for any semiclassical initial conditions, that is, for $(q(t_0), \cdots, \dddot{q}(t_0))$ satisfying $\dot{q}(t_0) \neq 0$, $|\ddot{q}(t_0)| << m|\dot{q}(t_0)|^3/(\hbar\sqrt{-2c_2})$ and $|\dddot{q}(t_0)| << m^2|\dot{q}(t_0)|^5/(-2\hbar^2 c_2)$, the solution of (2.20) is written as [f]

$$q(t) = vt + \sum_{n=1}^{\infty}(a_n\cos n\omega t + b_n\sin n\omega t) = vt + \sum_{n=1}^{\infty}A_n\cos(n\omega t + \theta_n) =: vt + f \ , \tag{2.26}$$

where $|v| = const. > 0$ and $|\dot{f}/v| << 1$; an additive const. was set to zero. The oscillating part is written as a Fourier series (Appendix E). The infinitesimal (inf.) order of A_n is $(\dot{f}/v)^n$ (Appendix E). The v, a_1, b_1 (A_1, θ_1) and the omitted additive const. work as initial conditions of the fourth order E-L eq. The ω is determined to $\omega = mv^2/(\hbar\sqrt{-2c_2}\,)$ from the E-L eq. truncated at $(\dot{f}/v)^1$. The $(A_2,\theta_2), (A_3,\theta_3), \cdots$ are determined recursively from (A_1,θ_1) and v. For example,

$$A_2\cos(2\omega t + \theta_2) = \frac{1}{2}\sqrt{1 - \frac{c_3^2}{8c_2^3}}\,\frac{\omega}{v}A_1^2\cos(2\omega t + 2\theta_1 + \xi) \ , \tag{2.27}$$

---

[f] We warn the reader that the ω in (2.26) is not the ω appearing in QM as the angular frequency of a massive particle, say, in a formula: $E = p^2/2m = \hbar\omega$. Let the ω representing the angular frequency of the massive particle be ω_Q, the ω in (2.26) be ω_L. In non-relativistic QM, from $p = mv = \hbar k$ and $v = dw_Q/dk$, we have $E = p^2/2m = \hbar\omega_Q$. While in our formulation, approximately, although $p = mv = \hbar k$ is common, v is expressed as $v = \omega_L/k$. We therefore have $E = p^2/2m = \hbar\omega_L/2$ (see (F.14)).

$$A_3\cos(3\omega t+\theta_3)=\frac{-3c_3}{2(-2c_2)^{\frac{3}{2}}}\frac{\omega A_1A_2}{v}\cos(3\omega t+\theta_1+\theta_2)-\frac{7}{6}\frac{\omega A_1A_2}{v}\sin(3\omega t+\theta_1+\theta_2)$$
$$+(\frac{5}{24}+\frac{c_4}{32c_2^2})(\frac{\omega A_1}{v})^2A_1\cos(3\omega t+3\theta_1)-\frac{7c_3}{12(-2c_2)^{\frac{3}{2}}}(\frac{\omega A_1}{v})^2A_1\sin(3\omega t+3\theta_1)\ ,$$

where $\omega=mv^2/(\hbar\sqrt{-2c_2}\,)$; the ξ is the angle satisfying $\cos\xi=\frac{-c_3}{(-2c_2)^{3/2}}/\sqrt{1-c_3^2/(8c_2^3)}$ and $\sin\xi=1/\sqrt{1-c_3^2/(8c_2^3)}$.

If the initial conditions are not of semiclassical, the solution is not necessarily wavy linear. Assume that initial conditions $(q(t_0),\cdots,\dddot{q}(t_0))$ satisfy $\dot{q}(t_0)>0$ and $p_{(1)}-\surd(p_{(1)}^2-2mE)<0$. (Note, if initial conditions is semiclassical, $p_{(1)}-\surd(p_{(1)}^2-2mE)\simeq p_{(1)}>0$ and $(p_{(1)}-\surd(p_{(1)}^2-2mE))/m$ is the minimum velocity (Appendix E).) Then, for $t\to+\infty$, the $\dot{q}$ asymptotically approaches some $const.>0$ because $\dot{q}$ cannot cross $\dot{q}=0$ (Appendix E).

The Lagrangian (2.19) in which at least one of c_1 and c_{2a} is not zero and the one in which $c_1=c_{2a}=0$ describe different systems in the sense that the relation between k and ω is different: $\omega=2kv$ for the former, $\omega=kv$ for the latter, which we see comparing p_{QHJ} and p_{EL} at the lowest inf. order of oscillation after inserting (2.26) into x of (2.16) and q of (2.23). In the present paper, we study the latter because we consider the latter is likely to describe the nature for reasons below. i) If $c_1\neq0$ (in general, if $c_n\neq0$, where n is odd) the Lagrangian is not invariant under time reversal. ii) If we require our free 3-D Lagrangian to be rotationally invariant as in CM, it is natural to consider that $c_1=c_{2a}=0$ (in general, $c_1=c_{2a}=c_3=c_5=\cdots=0$) because the c_1- and c_{2a}-terms (in general, c_1-, c_{2a}-, c_3-, c_5-, $\cdots$-terms) of (2.19) has no natural such 3-D form, which is contrasted to other terms having such 3-D forms with $\dot{q}^2\to\dot{\mathbf{q}}\cdot\dot{\mathbf{q}}=q_1^2+q_2^2+q_3^3$, etc. [g] iii) If $c_{2a}\neq0$, the Lagrangian is of the third order. However, because the fourth order E-L eq. is obtained from the simpler second order Lagrangian, we have no reason to adopt the third order Lagrangian.

We determine c_{2b} $(c_1=c_{2a}=0)$ comparing p_{QHJ} and p_{EL}. For the present purpose, we write p_{QHJ} or (2.16) in the form:

$$p_{QHJ}=\frac{\hbar k(1+\varepsilon_1)}{(\sin kx+\varepsilon_2\cos kx)^2+(1+\varepsilon_1)^2\cos^2 kx}\ . \tag{2.28}$$

In accordance with the present semiclassical setting, we replaced $l_1\to-(1+\varepsilon_1)$ and $l_2\to\varepsilon_2$, where $\varepsilon_1,\varepsilon_2<<1$. Inserting $q(t)$ of (2.26) into q of p_{EL} (2.23) and x of p_{QHJ} (2.28), and comparing p_{QHJ} and p_{EL} at the lowest inf. order of oscillation or at ε^1, where $\varepsilon=\sqrt{\varepsilon_1^2+\varepsilon_2^2}$, we have c_{2b} together with a_1,b_1, ω and v as the function of ε_1, ε_2, and E (Appendix F). The results are: $\omega=kv$, $v=\sqrt{2mE}(1-\varepsilon+\mathcal{O}(\varepsilon^2))/m$, $c_{2b}=-(1-2\varepsilon+\mathcal{O}(\varepsilon^2))/2$, where $v>0$ is assumed. The oscillation amplitude (a_1,b_1) are given as functions of ε_1, ε_2, and E in (F.16), (F.17). The inf. order of $\dot{f}/v$ with respect to ε is $1/2$:

---

[g] It seems that we can construct rotationally invariant 3-D forms of the $c_1,c_{2a},c_3,\cdots$-terms of (2.19) if we take the absolute value of them. This idea does not work because the resulting E-L eq. is unacceptable. For example, the E-L eq. obtained from $L=m\dot{q}^2/2+c_{2a}\hbar^2|\dddot{q}|/m|\dot{q}|^3$ is $m\ddot{q}\mp(3c_{2a}\hbar^2/m)(2\ddddot{q}/\dot{q}^4-16\ddot{q}\dddot{q}/\dot{q}^5+20\ddot{q}^3/\dot{q}^6)=0$, where '$-$'-sign for ($\dot{q}>0$, $\dddot{q}\geq0$), '$+$'-sign for ($\dot{q}>0$, $\dddot{q}<0$), which does not give a wavy linear trajectory as is seen from the truncated eq.: $m\ddot{q}\mp6c_{2a}\hbar^2\ddddot{q}/m\dot{q}^4=0$.

$\dot{f}/v \in \mathcal{O}(\varepsilon^{1/2})$. The 1-D free Lagrangian is written as

$$L = \frac{1}{2}m\dot{q}^2 - \frac{\hbar^2(1-2\varepsilon+\mathcal{O}(\varepsilon^2))}{2m}\frac{\ddot{q}^2}{\dot{q}^4} + \mathcal{O}(\hbar^3) \ . \tag{2.29}$$

The c_{2b} is fixed not only by the $(\varepsilon_1, \varepsilon_2, E)$ but also by the $(\dot{q}(t_0), \ddot{q}(t_0), \dddot{q}(t_0))$ or by the (v, a_1, b_1) of (2.26) since they are related through (F.13), (F.16), (F.17), and (2.26). The dependence of c_{2b} on $(\dot{q}(t_0), \ddot{q}(t_0), \dddot{q}(t_0))$ or on (v, a_1, b_1) is admissible (Appendix D).

No value of c_3 makes $p_{QHJ} = p_{EL}$ at the inf. order $\varepsilon^{3/2}$ of p_{QHJ} (Appendix G), from which we conclude that there is no Lagrangian beyond the semiclassical region, at least within our theoretical framework.

From (2.29), omitting $\mathcal{O}(\varepsilon^2)$ and noting $\varepsilon = const.$, we obtain

$$\text{(E-L eq.)} \quad 0 = m\ddot{q} + \frac{\hbar^2(1-2\varepsilon)}{m}\Big(\frac{\ddddot{q}}{\dot{q}^4} - \frac{8\dddot{q}\ddot{q}}{\dot{q}^5} + \frac{10\ddot{q}^3}{\dot{q}^6}\Big) + \mathcal{O}(\hbar^4) \tag{2.30a}$$

$$p_{(1)} = m\dot{q} - \frac{\hbar^2(1-2\varepsilon)}{m}\Big(\frac{2\ddot{q}^2}{\dot{q}^5} - \frac{\dddot{q}}{\dot{q}^4}\Big) + \mathcal{O}(\varepsilon^{\frac{3}{2}}) \ , \ \ p_{(2)} = -\frac{\hbar^2(1-2\varepsilon)}{m}\frac{\ddot{q}}{\dot{q}^4} + \mathcal{O}(\varepsilon^{\frac{3}{2}}) \tag{2.30b}$$

$$E = \frac{1}{2}m\dot{q}^2 - \frac{\hbar^2(1-2\varepsilon)}{m}\Big(\frac{5\ddot{q}^2}{2\dot{q}^4} - \frac{\dddot{q}}{\dot{q}^3}\Big) + \mathcal{O}(\varepsilon^{\frac{3}{2}}) \tag{2.30c}$$

$$p_{EL} = p_{(1)} + p_{(2)}\frac{\ddot{q}}{\dot{q}} = m\dot{q} - \frac{\hbar^2(1-2\varepsilon)}{m}\Big(\frac{3\ddot{q}^2}{\dot{q}^5} - \frac{\dddot{q}}{\dot{q}^4}\Big) + \mathcal{O}(\varepsilon^{\frac{3}{2}}) \ , \tag{2.30d}$$

where we have set c_3 hidden in $\mathcal{O}(\hbar^3)$ of (2.29) to zero considering the Lagrangian is invariant under time reversal. A benefit of which is that we have $\mathcal{O}(\varepsilon^{3/2})$ rather than $\mathcal{O}(\varepsilon^1)$ in (2.30b) and (2.30c). Note $\mathcal{O}(\varepsilon^{3/2}) = \mathcal{O}((\dot{f}/v)^3)$.

Next, we determine the semiclassical Lagrangian of a 3-D free particle corresponding to a 3-D momentum:

$$p_{QHJ\,i}(\mathbf{x}) = \frac{\hbar k_i(1+\varepsilon_{1i})}{(\sin\mathbf{kx} + \varepsilon_{2i}\cos\mathbf{kx})^2 + (1+\varepsilon_{1i})^2\cos^2\mathbf{kx}} \ , \tag{2.31}$$

where $i = 1,2,3$. We assume, from (2.19), the 3-D Lagrangian takes the form: $L = m\dot{\mathbf{q}}^2/2 + c_{2b}^{3D}\hbar^2\ddot{\mathbf{q}}^2/m\dot{\mathbf{q}}^4 + \mathcal{O}(\hbar^4)$, where $\dot{\mathbf{q}} = (\,\dot{q}_1(t), \dot{q}_2(t), \dot{q}_3(t)\,)$, $\dot{\mathbf{q}}^2 = \dot{\mathbf{q}}\cdot\dot{\mathbf{q}} = \dot{q}_1\dot{q}_1 + \dot{q}_2\dot{q}_2 + \dot{q}_3\dot{q}_3$, and $\dot{\mathbf{q}}^4 = (\dot{\mathbf{q}}\cdot\dot{\mathbf{q}})^2$; likewise for $\ddot{\mathbf{q}}^2$. We set c_3^{3D} corresponding to c_3 to zero ($c_3^{3D} = 0$) because there is no natural 3-D form corresponding to the c_3-term (footnote g). Then, in a similar manner as in the 1-D case, the c_{2b}^{3D} is determined as the function of $(\varepsilon_{1i}, \varepsilon_{2i}, E_i)$ in the form:

$$c_{2b}^{3D} = -\frac{m^2\mathbf{v}^4}{2\hbar^2\omega^2} = \frac{-m^2(v_1^2+v_2^2+v_3^2)^2}{2(v_1\sqrt{2mE_1} + v_2\sqrt{2mE_2} + v_3\sqrt{2mE_3})^2}$$
$$= -\frac{1}{2}\Big(1 - \frac{2\sum_{i=1}^{3}E_i\sqrt{\varepsilon_{1i}^2+\varepsilon_{2i}^2}}{E}\Big) + \mathcal{O}(\boldsymbol{\varepsilon}^{\frac{3}{2}}) \ ,$$

where $\omega = \sum_{i=1}^{3} k_i v_i$, $k_i = \sqrt{2mE_i}/\hbar$, $v_i \simeq (\sqrt{2mE_i}/m)(1 - \sqrt{\varepsilon_{1i}^2 + \varepsilon_{2i}^2}\ )$, $\boldsymbol{\varepsilon} = \sum_{i=1}^{3}\sqrt{}(\varepsilon_{1i}^2 + \varepsilon_{2i}^2)$ (Appendix H).

The semiclassical Lagrangian of a 3-D particle in a slowly changing potential field $V(\mathbf{q})$ is naturally considered to be

$$L = \frac{1}{2}m\dot{\mathbf{q}}^2 - \frac{\hbar^2(1-\tilde{\boldsymbol{\varepsilon}}(\mathbf{q}))}{2m}\frac{\ddot{\mathbf{q}}^2}{\dot{\mathbf{q}}^4} - V(\mathbf{q}) + \mathcal{O}(\hbar^3) \;, \tag{2.32}$$

where $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q}) << 1$, which is left undetermined, is a slowly changing function.

The Lagrangian (2.29) and (2.32) is regular, that is the Hessian does not vanish ($\partial^2 L/\partial\ddot{q}^2 \neq 0$ for (2.29), $|\partial^2 L/\partial\ddot{q}^i\partial\ddot{q}^j| \neq 0$ for (2.32)), a consequence of which is that the corresponding E-L eq. is fourth order and determines a unique vector field on the third-order tangent bundle T^3M. [6]

The 3-D free Lagrangian, or (2.32) with $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q}) = const.$ and $V(\mathbf{q}) = 0$, is translationally and rotationally invariant. The J-O momentum $\mathbf{p}_{(1)}$ is the Noether current asocciated to the invariance of the Lagrangian under spatial translation. [5]

From (2.32), ignoring $\mathcal{O}(\hbar^3)$ and $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q}) << 1$ (if interference is involved, we cannot ignore $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q})$; see the paragraph after next), we obtain (Appendix C)

$$\begin{aligned}\text{(E-L eq.)}\;\; 0 = -\frac{\partial V(\mathbf{q})}{\partial q_i} - m\ddot{q}_i + \frac{\hbar^2}{m}\Big(&-\frac{\ddddot{q}_i}{\dot{\mathbf{q}}^4} - \frac{4\dot{q}_i\dddot{\mathbf{q}}\ddot{\mathbf{q}}}{\dot{\mathbf{q}}^6} + \frac{8\dddot{q}_i\dot{\mathbf{q}}\ddot{\mathbf{q}}}{\dot{\mathbf{q}}^6} + \frac{4\ddot{q}_i\dot{\mathbf{q}}\dddot{\mathbf{q}}}{\dot{\mathbf{q}}^6} + \frac{2\ddot{q}_i\ddot{\mathbf{q}}\ddot{\mathbf{q}}}{\dot{\mathbf{q}}^6}\\ &+ \frac{12\dot{q}_i(\ddot{\mathbf{q}}\ddot{\mathbf{q}})(\dot{\mathbf{q}}\ddot{\mathbf{q}})}{\dot{\mathbf{q}}^8} - \frac{24\ddot{q}_i(\dot{\mathbf{q}}\ddot{\mathbf{q}})^2}{\dot{\mathbf{q}}^8}\Big)\end{aligned} \tag{2.33}$$

$$p_{(1)i} = \frac{\partial L}{\partial\dot{q}_i} - \frac{d}{dt}\frac{\partial L}{\partial\ddot{q}_i} = m\dot{q}_i + \frac{\hbar^2}{m}\Big(\frac{\dddot{q}_i}{\dot{\mathbf{q}}^4} + \frac{2\ddot{\mathbf{q}}^2\dot{q}_i}{\dot{\mathbf{q}}^6} - \frac{4\dot{\mathbf{q}}\ddot{\mathbf{q}}\ddot{q}_i}{\dot{\mathbf{q}}^6}\Big) \;,\;\; p_{(2)i} = -\frac{\hbar^2}{m}\frac{\ddot{q}_i}{\dot{\mathbf{q}}^4} \tag{2.34}$$

$$E = \mathbf{p}_{(1)}\dot{\mathbf{q}} + \mathbf{p}_{(2)}\ddot{\mathbf{q}} - L = \frac{1}{2}m\dot{\mathbf{q}}^2 + \frac{\hbar^2}{m}\Big(\frac{\dot{\mathbf{q}}\dddot{\mathbf{q}}}{\dot{\mathbf{q}}^4} + \frac{3}{2}\frac{\ddot{\mathbf{q}}^2}{\dot{\mathbf{q}}^4} - \frac{4(\dot{\mathbf{q}}\ddot{\mathbf{q}})^2}{\dot{\mathbf{q}}^6}\Big) + V(\mathbf{q}) \;. \tag{2.35}$$

The 3-D momentum $\mathbf{p} = \nabla\mathcal{S}_0$ is obtained as follows. From $\mathcal{S} = \int L dt$ and $E = \mathbf{p}_{(1)}\dot{\mathbf{q}} + \mathbf{p}_{(2)}\ddot{\mathbf{q}} - L$, we see, along the solution path, $d\mathcal{S} = d\mathcal{S}_0 - E dt = L dt = (\mathbf{p}_{(1)}\dot{\mathbf{q}} + \mathbf{p}_{(2)}\ddot{\mathbf{q}} - E)dt = (p_{(1)1} + p_{(2)1}\, \ddot{q}_1/\dot{q}_1)dq_1 + (\cdots)dq_2 + (p_{(1)3} + p_{(2)3}\, \ddot{q}_3/\dot{q}_3)dq_3 - E dt$. We therefore have

$$\frac{\partial\mathcal{S}_0}{\partial x_i} = p_{(1)i} + p_{(2)i}\frac{\ddot{q}_i}{\dot{q}_i} = m\dot{q}_i + \frac{\hbar^2}{m}\Big(\frac{\dddot{q}_i}{\dot{\mathbf{q}}^4} + \frac{2\ddot{\mathbf{q}}^2\dot{q}_i}{\dot{\mathbf{q}}^6} - \frac{4\dot{\mathbf{q}}\ddot{\mathbf{q}}\ddot{q}_i}{\dot{\mathbf{q}}^6}\Big) - \frac{\hbar^2}{m}\frac{\ddot{q}_i^2}{\dot{\mathbf{q}}^4\dot{q}_i}\,\Big|_{\mathbf{q}=\mathbf{x}} \;. \tag{2.36}$$

Although variables of 3-D free QSHJE separate on the Cartesian coordinates, those of the 3-D free E-L eq., or (2.33) with $V(\mathbf{q}) = 0$, does not, which however does not mean that two formalisms are inconsistent. The criterion of the consistency is whether physical quantities defined in both give the same values in both. In the present case, those defined in both are E [h] and $\partial\mathcal{S}_0/\partial x_i$. They give the same values up to $\boldsymbol{\varepsilon}^1$. The two therefore are consistent up to $\boldsymbol{\varepsilon}^1$. We cannot discuss the consistency at higher orders of $\boldsymbol{\varepsilon}$ because the Lagrangian formalism is not effective there (p.12).

In the interference region, we cannot ignore $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q})$ of (2.32) if we use the E-L eq. to determine the trajectory. However it is difficult to determine $\tilde{\boldsymbol{\varepsilon}}(\mathbf{q})$. We overcome this problem

[h] The author does not know whether i-direction component E_i of E is defined in the Lagrangian formalism or not. At least, there is no formula to express E_i.

bypassing the E-L eq., or reinterpreting (2.36) as a trajectory determining eq. Originally, (2.36) gives $\partial\mathcal{S}_0/\partial x_i$ from a given $\mathbf{q}(t)$. However, since the $\partial\mathcal{S}_0/\partial x_i$ on the LHS is given also as a solution of the QSHJE, we regard (2.36) as a differential eq. for $\mathbf{q}(t)$ after inserting a solution of the QSHJE on the LHS. With this reinterpretation, we can determine the trajectory even in the interference region since the QSHJE, which is a partial differential eq., describes the interference; see §3.3.

3. Application

Based on the formalism developed in §2, we explain how quantum phenomena relevant to a spinless massive particle appear.

3.1. *Energy quantization in 1-D space, Uncertainty relation, Tunneling, and Free particle*

Energy quantization in 1-D space The energy of the QSHJE with a confining potential is quantized to the same levels as that of QM, which is seen from (2.12).

Uncertainty relation between position and momentum We examine a particle in a 1-D potential well. Let the energy E of a particle be $E > 0$. Let the potential of the well be $V(x) = 0$ for $|x| \leq L,\ V(x) >> E$ for $|x| > L$. Let $k = \sqrt{2mE}/\hbar$. We assume that the particle has the lowest possible energy. Then we can take $\Psi(x) \simeq \cos\frac{\pi}{2L}x$ and $\Psi^D(x) \simeq \sin\frac{\pi}{2L}x$ as two linearly independent solutions of the corresponding SE. The momentum $p = \partial\mathcal{S}_0/\partial x$ in the well is found using the formula (2.9). Assuming $l_2 = 0$, we obtain (note $W = \Psi'\Psi^D - \Psi^{D\prime}\Psi = -\frac{\pi}{2L}$)

$$p = \frac{\partial\mathcal{S}_0}{\partial x} = \frac{\hbar W l_1}{(\Psi^D + l_2\Psi)^2 + l_1^2\Psi^2} = \frac{-\frac{\hbar\pi l_1}{2L}}{\sin^2\frac{\pi x}{2L} + l_1^2\cos^2\frac{\pi x}{2L}} = \frac{-\frac{\hbar\pi l_1}{2L}}{1 + (l_1^2 - 1)\cos^2\frac{\pi x}{2L}} \ .$$

From which, assuming $l_1 > 0$, $\Delta p := \text{max.}|p| - \text{min.}|p|$ is obtained as

$$\Delta p = \frac{\hbar\pi}{\Delta x}\ |l_1 - \frac{1}{l_1}|\ ,$$

where the width $2L$ of the well was replaced with Δx. If $l_1 \simeq 1.2$ or 0.85, the quantum mechanical uncertainty relation $\Delta p\Delta x \simeq \hbar$ is recovered.

Tunneling From the energy formula (2.35), we see that it can be $\dot{\mathbf{q}}^2 > 0$ even in a region where $E - V(\mathbf{x}) < 0$, or that tunneling occurs respecting the energy conservation law and the particle picture. We note that we cannot determine the trajectory in the region where $E \approx V(\mathbf{x})$ since the semiclassicality assumption breaks down there.

A free particle The trajectory of a free particle is not necessarily linear: $\mathbf{q}(t) = \mathbf{v}t$ ($\mathbf{v} =$ *const*.) since our E-L eq. (2.33) is of the 4th order. The deviation from $\mathbf{q}(t) = \mathbf{v}t$, if any, is estimated to be very small, say, from Time-of-Flight experiments. When the deviation is small, the trajectory is approximated by $\mathbf{q}(t) = \mathbf{v}t + \boldsymbol{\alpha}\cos(\omega t + \boldsymbol{\theta})$. Inserting this $\mathbf{q}(t)$ into the energy formula (2.35), we see that the energy E of the particle is expressed as

$$E \simeq \frac{1}{2}m\mathbf{v}^2 + \frac{1}{2}m\boldsymbol{\alpha}^2\omega^2\ . \tag{3.1}$$

Even if the particle is charged, the electromagnetic radiation does not occur. Such radiation to occur, it has to be $m\omega^2\boldsymbol{\alpha}^2/2 \geq \hbar\omega$. Substituting $\hbar\omega$ with $m\mathbf{v}^2$ (see footnote f), we see that $\omega^2\boldsymbol{\alpha}^2/2\mathbf{v}^2 \geq 1$, which contradicts the semiclassicality assumption in which the oscillation is small.

3.2. *Energy quantization in 3-D space and angular momentum quantization*

Phenomena corresponding to the quantizations of l_z (z-component of the angular momentum), l^2 (square of the angular momentum), and E (energy) of QM appear to the system of a particle in a 3-D attractive central force field $V(r)$.

The QSHJE of which is separated to three 1-D eqs. on the spherical polar coordinate system connected to the Cartesian one with $x = r\sin\theta\cos\varphi$, $y = r\sin\theta\sin\varphi$, $z = r\cos\theta$. That is, the QSHJE

$$\frac{\hbar^2}{2m}\frac{\nabla^2 R}{R} - \frac{1}{2m}(\nabla\mathcal{S}_0)^2 - V(r) + E = 0\ , \qquad \nabla(\frac{R^2}{m}\nabla\mathcal{S}_0) = 0 \tag{3.2}$$

is, setting $R = R_\varphi R_\theta R_r$, $\mathcal{S}_0 = \mathcal{S}_{0\varphi} + \mathcal{S}_{0\theta} + \mathcal{S}_{0r}$, and making use of the technique used to derive (2.6) from (2.5), separated to

$$-\frac{\hbar^2}{4m}\{e^{\frac{2i}{\hbar}\mathcal{S}_{0\varphi}}, \varphi\} = -\frac{\hbar^2}{2m}\mathrm{m}^2\ , \quad -\frac{\hbar^2}{4m}\{e^{\frac{2i}{\hbar}\mathcal{S}_{0\theta}}, \theta\} = -\frac{\hbar^2}{2m}\Big(\lambda - \frac{\mathrm{m}^2}{\sin^2\theta} + \frac{1}{2} + \frac{\cos^2\theta}{4\sin^2\theta}\Big)\ ,$$

$$-\frac{\hbar^2}{4m}\{e^{\frac{2i}{\hbar}\mathcal{S}_{0r}}, r\} = -\frac{\hbar^2}{2m}\Big(\frac{2m}{\hbar^2}(E - V(r)) - \frac{\lambda}{r^2}\Big)\ , \tag{3.3}$$

where $\frac{\hbar^2}{2m}\mathrm{m}^2$, $\frac{\hbar^2}{2m}\lambda$ are separation consts. in the first eq. of (3.2). The separation consts. in the second eq. of (3.2) were set to zeros.

First, we look for requirements for the solution of (3.2) to be physically acceptable. **1**) In 1-D space, the solution $\mathcal{S}_0(q)$ of the QSHJE with a confining potential is required to satisfy (2.12). In 3-D space, if QSHJE is separable on the Cartesian coordinates to three 1-D eqs., we can express the 3-D version of (2.12), requiring the (2.12) on each direction, as 'The corresponding 3-D SE has a L^2-solution.' $\cdots(\star)$ Since in this form, the requirement is independent from coordinate systems, we impose this requirement on the 3-D QSHJE with a confining potential regardless of the separability. **2**) In 1-D space, the solution $\mathcal{S}_0(x)$ has to be a diffeomorphism from x to $\mathcal{S}_0$. In 3-D space, this requirement is translated to: $\nabla\mathcal{S}_0 \neq 0, \pm\infty$ except at points at which the effective potential $V_{\text{eff}} = \hbar^2(4\mathrm{m}^2 - \cos^2\theta)/(8m\sin^2\theta)$ or $V_{\text{eff}} = V(r) + \hbar^2\lambda/(2mr^2)$ is singular ($V_{\text{eff}} = \pm\infty$). **3**) In φ-direction, we require that i) $\partial\mathcal{S}_{0\varphi}/\partial\varphi$ is single-valued, ii) $\int_0^\pi \partial\mathcal{S}_{0\varphi}/\partial\varphi d\varphi = nh/2$ ($n = \pm1, \pm2, \cdots$), which is understood as follows. Consider the 3-D QSHJE having the confining potential separable on both Cartesian and spherical polar coordinates, that is the QSHJE with the harmonic potential. Then, on each of x-, y-, and z-directions, the (2.11) is imposed on the solution $\mathcal{S}_{0i}$ ($i = x, y, z$). Since $\mathcal{S}_0(x, y, z)$ is a scaler, the integral value of the (2.11) on each direction is independent from the integral path taken in 3-D space. Taking the integral path of (2.11) for $\mathcal{S}_{0x}$ along the arc: $r = const. >> 1$, $\theta = \pi/2$, $\pi \leq \varphi \leq 2\pi$, we see that the above requirement is equivalent to the (2.11) for $\mathcal{S}_{0x}$. If $\partial\mathcal{S}_{0\theta}/\partial\theta \neq 0$ or $\partial\mathcal{S}_{0r}/\partial r \neq 0$, $n = 0$ or $\partial\mathcal{S}_{0\varphi}/\partial\varphi \equiv 0$ is

allowed since $\nabla \mathcal{S}_0 \neq 0$ for which. Summarizing; For a solution $\mathcal{S}_0$ of the 3-D QSHJE with an attractive central force field to be physically acceptable,
$\star$1) The 3-D SE corresponding to the 3-D QSHJE has a L^2-solution.
$\star$2) $\nabla \mathcal{S}_0 \neq 0, \pm\infty$ except at singular points of effective potential.
$\star$3) $\partial \mathcal{S}_{0\varphi}/\partial\varphi$ is single-valued.
$\star$4) $\int_0^\pi \partial \mathcal{S}_{0\varphi}/\partial\varphi d\varphi = nh/2$ ($n = \pm 1, \pm 2, \cdots$). If $\partial \mathcal{S}_{0\theta}/\partial\theta \neq 0$ or $\partial \mathcal{S}_{0r}/\partial r \neq 0$, $n = 0$ is allowed.

Next, we examine for which values of m, λ and E, the $\star 1) \sim \star 4)$ are satisfied. The solutions of (3.3) are constructed from the solutions of the 1-D SE's corresponding to (3.3), which are obtained separating the 3-D SE corresponding to (3.2):

$$-\frac{\hbar^2}{2m}\nabla^2\psi + (V(r) - E)\psi = 0 \tag{3.4}$$

with $\psi(\varphi,\theta,r) = \Phi(\varphi)\Theta(\theta)\mathsf{R}(r)$. Explicitly, they are

$$\frac{d^2\Phi}{d\varphi^2} + \mathsf{m}^2\Phi = 0\ , \tag{3.5a}$$

$$\frac{1}{\sin\theta}\frac{d}{d\theta}(\sin\theta\frac{d\Theta}{d\theta}) + (\lambda - \frac{\mathsf{m}^2}{\sin^2\theta})\Theta = 0\ , \tag{3.5b}$$

$$\frac{1}{r}\frac{d^2(r\mathsf{R})}{dr^2} + (\frac{2m}{\hbar^2}(E - V) - \frac{\lambda}{r^2})\mathsf{R} = 0\ . \tag{3.5c}$$

Allowed values for m If $\mathsf{m} \neq 0$, the general solution of (3.5a) is obtained from two linearly independent solutions $\Phi_1 = \cos\mathsf{m}\varphi$ and $\Phi_2 = \sin\mathsf{m}\varphi$ as $\Phi = c_1\cos\mathsf{m}\varphi + c_2\sin\mathsf{m}\varphi$, where $c_1, c_2 \in \mathbb{C}$. The $\partial \mathcal{S}_{0\varphi}/\partial\varphi$ is constructed from which, using (2.15), as

$$\frac{\partial \mathcal{S}_{0\varphi}}{\partial\varphi} = \frac{\hbar}{i}\frac{\mathsf{m}(\bar{c}_1 c_2 - c_1\bar{c}_2)}{(c_1\bar{c}_1 + c_2\bar{c}_2) + (c_1\bar{c}_1 - c_2\bar{c}_2)\cos 2\mathsf{m}\varphi + (c_1\bar{c}_2 + c_2\bar{c}_1)\sin 2\mathsf{m}\varphi} \in \mathbb{R}\ .$$

Helped by the formula: $\int \frac{dx}{(a\sin x + b\cos x + c)} = \frac{2}{\sqrt{c^2-a^2-b^2}}\arctan\left(\frac{(c-b)\tan(x/2)+a}{\sqrt{c^2-a^2-b^2}}\right)$, where $c^2 > a^2 + b^2$, we see that, to make $\int_0^\pi \partial \mathcal{S}_{0\varphi}/\partial\varphi d\varphi = nh/2$ true, it has to be $\mathsf{m} = \pm 1, \pm 2, \cdots$.

If $\mathsf{m} = 0$, we take $\Phi_1 = \mathbf{1}$ (see p.7) and $\Phi_2 = \varphi$ as two linearly independent solutions of (3.5a). From which the general solution is obtained as $\Phi = c_1\mathbf{1} + c_2\varphi$. The $\partial \mathcal{S}_{0\varphi}/\partial\varphi$ is constructed using (2.15) as

$$\frac{\partial \mathcal{S}_{0\varphi}}{\partial\varphi} = \frac{\hbar}{2i}\frac{(\bar{c}_1 c_2 - c_1\bar{c}_2)\mathbf{1}}{(\ c_1\bar{c}_1\mathbf{1}^2 + (c_1\bar{c}_2 + c_2\bar{c}_1)\mathbf{1}\varphi + c_2\bar{c}_2\varphi^2\ )} \in \mathbb{R}\ .$$

If $\bar{c}_1 c_2 \neq c_1\bar{c}_2$, $\mathcal{S}_{0\varphi}$ does not satisfy the $\star$3). If $\bar{c}_1 c_2 = c_1\bar{c}_2$, $\mathcal{S}_{0\varphi}$ satisfies the $\star$3) with $\partial \mathcal{S}_{0\varphi}/\partial\varphi \equiv 0$, which is allowed because θ-direction momentum can be non-zero as is seen below (see $\star$2)). Thus, allowed values for m are $\mathsf{m} = 0, \pm 1, \pm 2, \cdots$.

Allowed values for λ and E Since the values of λ and E for which the $\star$1) is satisfied are found in standard textbooks of QM, we do not repeat the derivation of them (allowed values for λ are $\lambda = l(l+1)$ ($l = 0, 1, \cdots$), those for E depend on $V(r)$). What we need to confirm is that the $\star$2) is satisfied for such λ's and E's. We have seen $\partial \mathcal{S}_{0\varphi}/\partial\varphi = 0$ for $\mathsf{m} = 0$. To satisfy the $\star$2), therefore, either of $\partial \mathcal{S}_{0\theta}/\partial\theta$ or $\partial \mathcal{S}_{0r}/\partial r$ has to be non-zero when

$\mathsf{m} = 0$. We will see that, for any physically acceptable values of m, λ, and E, $\partial\mathcal{S}_{0r}/\partial r \neq 0$ is not allowed, whereas $\partial\mathcal{S}_{0\theta}/\partial\theta \neq 0$ is allowed.

For values of λ and E which give a L^2-solutions for (3.4), the general solution of (3.5b) and (3.5c) are written as $\Theta = c_3\Theta_{\mathrm{con}} + c_4\Theta_{\mathrm{div}}$ and $\mathsf{R} = c_5\mathsf{R}_{\mathrm{con}} + c_6\mathsf{R}_{\mathrm{div}}$, where $\Theta_{\mathrm{con}} \cdots \mathsf{R}_{\mathrm{div}} \in \mathbb{R}$; the Θ_{con} ($\mathsf{R}_{\mathrm{con}}$) is the θ- (r-) direction component of the L^2-solution; the Θ_{div} ($\mathsf{R}_{\mathrm{div}}$) is the solution linearly independent from Θ_{con} ($\mathsf{R}_{\mathrm{con}}$). For the $\partial\mathcal{S}_{0\theta}/\partial\theta$ ($\partial\mathcal{S}_{0r}/\partial r$) to be non-zero, both of c_3 and c_4 (c_5 and c_6) has to be non-zero (see (2.15)). However, $c_4(c_6) \neq 0$ is not necessarily allowed because 3-D SE (3.4) is not necessarily satisfied by the solution $\psi = \Phi\Theta\mathsf{R}$ constructed from the 1-D solutions. Since $r = 0$ and $\theta = 0, \pi$ are singular points as is seen from the Jacobian J: $J = r^2 \sin\theta$, the behavior of $\psi = \Phi\Theta\mathsf{R}$ there may be incompatible with the 3-D SE on the Cartesian coordinate system. We therefore examine the behavior of $\psi = \Phi\Theta\mathsf{R}$ there.

As a preparation, we give the general solution of (3.5b) around $\theta = 0$ for $\mathsf{m} = 0$ and $\lambda \in \mathbb{C}$. It is obtained, using the Frobenius method (e.g., Ref. 16), as [i]

$$\Theta(\cos\theta) = G\Theta_1(\cos\theta) + H\Theta_2(\cos\theta) = GF(-l, l+1; 1; \frac{1-\cos\theta}{2})$$
$$+ H\sum_{r=0}^{\infty} \frac{(-l)_r(l+1)_r}{(1)_r^2} \Big(\ln\frac{1-\cos\theta}{2} + \sum_{k=0}^{r-1}\Big(\frac{1}{-l+k} + \frac{1}{l+1+k} - \frac{2}{1+k}\Big)\Big)\Big(\frac{1-\cos\theta}{2}\Big)^r,$$

where $G, H = const. \in \mathbb{C}$; $l(l+1) = \lambda$; $(\bullet)_0 = 1$, $(\bullet)_r = \bullet\cdots(\bullet + r - 1)$ for $r \geq 1$; the $\sum_{k=0}^{-1}(\cdots)$ is understood as zero; the $F(\alpha, \beta; \gamma; z)$ is the hypergeometric function defined as $F(\alpha,\beta;\gamma;z) := \sum_{r=0}^{\infty}(\alpha)_r(\beta)_r z^r/((\gamma)_r r!)$ $(|z| < 1)$. Note the $1/(-l+k) = 1/0$ and $1/(l+1+k) = 1/0$ are cancelled away with a factor in the numerator of $\frac{(-l)_r(l+1)_r}{(1)_r^2}$. We know $F(-l, l+1; 1; \frac{1-\cos\theta}{2})$ is a L^2-function on $0 \leq \theta \leq \pi$ for $l \in \mathbb{Z}$. Since $l = 0, 1, \cdots$ and $l = -1, -2, \cdots$ give the same Θ, we restrict the domain of l to $l \geq 0$. To list a few general solutions, $\Theta(\cos\theta) = G + H(\ln\frac{1-\cos\theta}{2} - \ln\frac{1+\cos\theta}{2})$ for $l = 0$, $\Theta(\cos\theta) = G\cos\theta + H(\cos\theta\ln\frac{1-\cos\theta}{2} - \cos\theta\ln\frac{1+\cos\theta}{2} + 2(1-\cos\theta))$ for $l = 1$.

First, we show $\partial\mathcal{S}_{0r}/\partial r = 0$. [j] We assume that the potential takes the form: $V(r) = a_{-1}/r + a_0 + a_1 r + \cdots$, where $a_{-1}, a_0, \cdots = const. \in \mathbb{R}$. If $l = 0$, $\chi(r) := r\mathsf{R}(r)$ behaves like r or a $const. \neq 0$ for $r \to 0$, which is seen from the indicial eq. (e.g., Ref. 16). The $\chi(r)$ which behaves like the latter is unacceptable. Assume the solution behaves like a $const. = b_0 \neq 0$ for $r \to 0$. Then, R behaves like b_0/r for $r \to 0$. The $\Psi = \Phi\Theta\mathsf{R} = (1+\Theta_{\mathrm{div}})b_0/r$ (we ignore irrelevant coefficients; the explicit form of Θ_{div} is given above) constructed with which is not a solution of the 3-D SE (3.4) as is seen as follows. If we insert $\psi = (1+\Theta_{\mathrm{div}})b_0/r$ into (3.4) and integrate the eq. over a small sphere Ω of which center is the origin, we see the integral of the LHS equals $2\hbar^2 b_0/m$, or $\lim_{\Omega\to 0}\int(-\hbar^2\nabla^2\psi/2m + (V - E)\psi)d\Omega = 2\hbar^2 b_0/m$. Note $\int\nabla^2(1/r)d\Omega = -4\pi$; $\int\nabla^2(\Theta_{\mathrm{div}} b_0/r)d\Omega = 0$ since $\Theta_{\mathrm{div}}(\frac{\pi}{2} -$

---

[i] General solutions of (3.5b) for any $\mathsf{m}, \lambda \in \mathbb{C}$ are obtained from general solutions of the hypergeometric eq.: $z(1-z)u'' + (\gamma - (\alpha+\beta+1))u' - \alpha\beta u = 0$ (e.g., Ref. 17). General solutions of which for any $\alpha, \beta, \gamma \in \mathbb{C}$ are found (exactly, were found on June 2010) in explicit forms with derivation on
http://en.wikipedia.org/wiki/Frobenius_solution_to_the_hypergeometric_equation.

[j] The following discussion is similar to that found in, e.g., chap.9 of Ref. 14. The difference is that ours involves Θ_{div} and $\mathsf{R}_{\mathrm{div}}$.

$\theta) = -\Theta_{\rm div}(\frac{\pi}{2}+\theta)$; $\lim_{\Omega\to 0}\int(V-E)\psi d\Omega = \lim_{\Omega\to 0}\int \frac{a_{-1}}{r}\frac{b_0}{r}(1+\Theta_{\rm div})r^2\sin\theta d\theta d\varphi = 0$. On the other hand, that of the RHS is zero. It therefore has to be $b_0 = 0$. Whereas the χ's which behave like r for $r \to 0$ are allowed. If $l \neq 0$, $\chi(r)$ behaves like r^{l+1} or r^{-l} for $r \to 0$. Solutions which behave like r^{-l} for $r \to 0$ are not allowed, which is is seen in a similar manner as above. Solutions which behave like r^{l+l} for $r \to 0$ are allowed. Thus, for any acceptable values of l, only one from the two linearly independent solutions: $\mathsf{R}_{\rm con}(r)$ and $\mathsf{R}_{\rm div}(r)$, is acceptable. (The $\chi(r)$'s which behave differently for $r \to 0$ are linearly independent.) Accordingly, $\partial\mathcal{S}_{0r}/\partial r = 0$.

Second, we show that $\partial\mathcal{S}_{0\theta}/\partial\theta \neq 0$ is allowed. To see $\psi = \Phi\Theta\mathsf{R}_{\rm con} = \Phi(c_3\Theta_{\rm con} + c_4\Theta_{\rm div})\mathsf{R}_{\rm con}$ with $c_4 \neq 0$ is allowed, we insert this ψ into (3.4) and integrate the eq. over small volume: $0 \leq \theta \leq \delta = const. << 1$ as in the case of the r-direction. Then, we see $\psi = \Phi(c_3\Theta_{\rm con} + c_4\Theta_{\rm div})\mathsf{R}_{\rm con}$ does not contradict (3.4), which is understood as a result of the slow divergence of $\Theta_{\rm div}$ for $\theta \to 0, \pi$. Thus, we see $\partial\mathcal{S}_{0\theta}/\partial\theta \neq 0$ is allowed.

3.3. *Interference*

We determine, in the setting of Fig.1, the trajectories of electrons in the interference (i.f.) region, and the density distribution of them on the screen. We calculate the trajectory with (2.36), the density distribution from the hitting positions. We will see that the density distribution differs from that of QM by a few percent.

• *Apparatus* We study with the apparatus of Fig.1. Explicit values are:
$k_x = \pm 4.99 \times 10^4 mm^{-1}, \quad k_y = 0, \quad k_z = 1.45 \times 10^9 mm^{-1},$
filament diameter: 4.15×10^{-4} mm,
aperture size: $-0.2mm \leq x \leq 0.2mm,\ -0.2mm \leq y \leq 0.2mm,$
the distance from the filament to the observation screen: 33.77mm,

where $k_x(k_y,k_z)$ is the wavenumber in the $x(y,z)$-direction after passing the electron biprism. The electron source is assumed to be located infinitely far from the biprism. With these values, we simulate the apparatus of Ref. 7

• *Incoming beam* In our formulation, only a superposed Gaussian beam (by 'beam', we mean the SE's solution) is allowed as a source side solution of the corresponding SE, which we call the incoming beam. The incoming beam is a solution of the free SE. The plane wave, the Bessel beam (e.g., Ref. 18) and the Gaussian beam (e.g., Ref. 19) are candidates. The superposition of different energy solutions is not allowed since the QSHJE, the corresponding SE too, has a fixed energy; see §4. From the plane wave, we cannot construct non-zero momentum in the x-direction behind the biprism (see (3.18)). The Bessel beam cannot be the incoming beam because the transverse amplitude of which changes sign as $|r - r_0|$, where r_0 is the beam center and r is the transverse distance from r_0, increases. We therefore take the Gaussian beam. The Gaussian beam, which is a solution of the paraxial form of the free SE, is written as

$$\Psi_G = \frac{w_0}{w(z)} \exp(-\frac{(x-x_0)^2+(y-y_0)^2}{w^2(z)}) \exp ik(z + \frac{(x-x_0)^2+(y-y_0)^2}{2\mathcal{R}(z)} + \frac{\hat{\phi}(z)}{k}) \ , \quad (3.6)$$

where w_0 is the beam width at the beam waist position, w is the beam width at z, (x_0, y_0) is the beam center coordinates, $k = \sqrt{(k_x^2 + k_y^2 + k_z^2)}$, $\mathcal{R}$ is the radius of curvature of the



Fig. 1. Experimental setting: Electrons are emitted from the source at the left side end and accelerated toward the screen. Electrons passed through the upper (lower) slot change the moving direction downward (upward) due to the voltage applied to the central filament.

wavefront, and $\hat{\phi}$ is the Gouy phase. The z-dependence of w, $\mathcal{R}$, and $\hat{\phi}$ is written using the Rayleigh range $z_R := kw_0^2/2$ as $w^2(z) = w_0^2(1+(z/z_R)^2)$, $\mathcal{R}(z) = z(1+(z_R/z)^2)$, and $\hat{\phi}(z) = \tan^{-1}(-z/z_R)$; see, e.g., Ref. 19 for details. We consider that the beam waist position, at which $\mathcal{R} = \infty$, is adjusted to $z = 0$ with focusing lenses on the source side. Because preliminary calculation suggests [k] that in the real world the transverse beam profile is close to $e^{-|r-r_0|}$ rather than $e^{-(r-r_0)^2}$, we introduce a superposed Gaussian beam. In the present calculation, we assume the simplest of which: two components Gaussian beam (see (3.15)). We assume, although we don't exclude other possibilities (see p.24), that the incoming beam accompanies an electron at its center. The conclusion that the diffraction intensity distribution differs from that of QM is kept intact regardless of the transverse electron distribution in the ensemble of identical incoming beams (see footnote s).

• *Momentum field behind the biprism* [l] We determine the momentum field behind the biprism caused by the Gaussian beam (3.6), from which the momentum field caused by the superposed Gaussian beam is easily obtained.

The momentum field $\mathbf{p}(\mathbf{x}) = \nabla \mathcal{S}_0(\mathbf{x})$ behind the biprism is constructed from the solution $\Psi_{\rm diff}$ of the corresponding SE using (2.15). The $\Psi_{\rm diff}$ is the superposition of two beams, one which originates from the virtual source located at the lower left far from the filament and was diffracted by the knife-edge open downward at $z = 0$ (no voltage applied), the other which is obvious from symmetry; see Fig.2 and Ref. 7. We use three coordinate systems (x,y,z), $(\tilde{x},\tilde{y},\tilde{z})$, and $(\tilde{\tilde{x}},\tilde{\tilde{y}},\tilde{\tilde{z}})$ which are related with



Fig. 2. The coordinate systems

$$\tilde{x} = \frac{k_z x - k_x z}{\sqrt{k_x^2+k_z^2}},\ \tilde{y} = y,\ \tilde{z} = \frac{k_x x + k_z z}{\sqrt{k_x^2+k_z^2}}, \quad \tilde{\tilde{x}} = \frac{k_z x + k_x z}{\sqrt{k_x^2+k_z^2}},\ \tilde{\tilde{y}} = y,\ \tilde{\tilde{z}} = \frac{-k_x x + k_z z}{\sqrt{k_x^2+k_z^2}}\ . \tag{3.7}$$

We consider two Gaussian beams $\tilde{\Psi}_G$ and $\tilde{\tilde{\Psi}}_G$. The $\tilde{\Psi}_G$ ($\tilde{\tilde{\Psi}}_G$) propagates parallel to the $\tilde{z}$ ($\tilde{\tilde{z}}$) axis toward the $\tilde{z}+$ ($\tilde{\tilde{z}}+$) direction. The $\tilde{\Psi}_G$ is diffracted by the knife-edge in the $(\tilde{x},\tilde{y})$ plane at $\tilde{z} = 0$ open downward from $\tilde{x} = -\mathrm{d}/2$ to $-\infty$ (d is the diameter of the filament). The $\tilde{\tilde{\Psi}}_G$ is diffracted by the knife-edge in the $(\tilde{\tilde{x}},\tilde{\tilde{y}})$ plane at $\tilde{\tilde{z}} = 0$ open upward from $\tilde{\tilde{x}} = +\mathrm{d}/2$ to $+\infty$.

---

[k] When the beam width w_0 is large enough to form a reasonable interference contrast on the screen, the electrons which passed close to the filament accumulate in a small area on the screen, which contradicts experiments.

[l] Although we find the formula which gives the Gaussian beam field diffracted by the knife-edge in Ref. 20, it is too elaborate for our purpose. We therefore determine the diffracted field in a simpler setting.

The disturbance $\tilde{U}(P)$ of $\tilde{\Psi}_G$ at the point $P = (\tilde{X}, \tilde{Y}, \tilde{Z})$ behind the downward open knife-edge, where $(\tilde{X}, \tilde{Y}, \tilde{Z})$ are coordinate values on the $(\tilde{x}, \tilde{y}, \tilde{z})$-system, is given as (e.g., Ref. 21 chap.8)

$$\tilde{U}(P) \simeq -\frac{iA}{\lambda \tilde{s}\,'} \int_{-\infty}^{-\mathsf{d}/2} d\tilde{\xi} \int_{-\infty}^{+\infty} d\tilde{\eta} \exp\Big(-\frac{(\tilde{\xi} - \tilde{x}_0)^2 + (\tilde{\eta} - \tilde{y}_0)^2}{w_0^2} + ik\tilde{s}\Big) \,, \tag{3.8}$$

where A: amplitude of the incoming beam; $\lambda (= 2\pi/k)$: wave length; $(\tilde{\xi}, \tilde{\eta})$: coordinates on the $(\tilde{x}, \tilde{y})$ plane at $\tilde{z} = 0$; $\tilde{s} = \sqrt{(\tilde{X} - \tilde{\xi})^2 + (\tilde{Y} - \tilde{\eta})^2 + \tilde{Z}^2}$, $\tilde{s}\,' = \sqrt{\tilde{X}^2 + \tilde{Y}^2 + \tilde{Z}^2}$. We consider only the case: $\tilde{y}_0 = 0$ and $\tilde{Y} = 0$. In this case, $\tilde{U}(P)$ of (3.8) is written, with the Fresnel approximation, as (see Appendix I)

$$\begin{aligned} \tilde{U}(P) = {} & \frac{Ae^{-i\pi/4}}{\sqrt{2(1-\tilde{l}^2)}} e^{ik\tilde{s}\,'} \\ & \times \Big((\cos(\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}}) \cdot \mathsf{M}_1 + \sin(\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}}) \cdot \mathsf{N}_1) + i(\cos(\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}}) \cdot \mathsf{N}_1 - \sin(\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}}) \cdot \mathsf{M}_1) \Big) \,. \end{aligned}$$

where $\tilde{l} = \tilde{X}/\tilde{s}\,'$, $\tilde{\alpha} = k(1 - \tilde{l}^2)/(\pi \tilde{s}\,')$, $\mathsf{M}_1 = \mathsf{M}_1(\tilde{X}, \tilde{Z})$ and $\mathsf{N}_1 = \mathsf{N}_1(\tilde{X}, \tilde{Z})$ are slowly changing (compared to $\cos 2k_x x$) functions of $\tilde{X}$ and $\tilde{Z}$, explicit forms of which are given in Appendix I. Rewriting $\tilde{U}(P)$ as $\tilde{U}(P) = \frac{A \exp i(k\tilde{s}\,' - \pi/4)}{\sqrt{2(1-\tilde{l}^2)}} (\mathsf{M}_1 + i\mathsf{N}_1) \exp(-i\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}})$, we see that the field intensity $|\tilde{U}(P)|^2$ caused by the lower-slot-passed beam is $\frac{A^2}{2}(\mathsf{M}_1^2 + \mathsf{N}_1^2)$ (note $\tilde{l}^{\,2} << 1$).

In a same manner, $\tilde{\tilde{U}}(P)$ is written as

$$\begin{aligned} \tilde{\tilde{U}}(P) = {} & \frac{Ae^{-i\pi/4}}{\sqrt{2(1-\tilde{\tilde{l}}^2)}} e^{ik\tilde{\tilde{s}}\,'} \\ & \times \Big((\cos(\frac{k^2 \tilde{\tilde{l}}^{\,2}}{2\pi\tilde{\tilde{\alpha}}}) \cdot \mathsf{M}_2 + \sin(\frac{k^2 \tilde{\tilde{l}}^{\,2}}{2\pi\tilde{\tilde{\alpha}}}) \cdot \mathsf{N}_2) + i(\cos(\frac{k^2 \tilde{\tilde{l}}^{\,2}}{2\pi\tilde{\tilde{\alpha}}}) \cdot \mathsf{N}_2 - \sin(\frac{k^2 \tilde{\tilde{l}}^{\,2}}{2\pi\tilde{\tilde{\alpha}}}) \cdot \mathsf{M}_2) \Big) \,, \end{aligned}$$

where what $\mathsf{M}_2, \mathsf{N}_2, \tilde{\tilde{l}}, \tilde{\tilde{\alpha}}$ represent are obvious.

The diffracted field Ψ_{diff} at $P = (X, Y, Z)$ behind the biprism is the superposition of $\tilde{U}(\tilde{X}, \tilde{Y}, \tilde{Z})$ and $\tilde{\tilde{U}}(\tilde{\tilde{X}}, \tilde{\tilde{Y}}, \tilde{\tilde{Z}})$:

$$\Psi_{\text{diff}} = \tilde{U}(P) + \tilde{\tilde{U}}(P) = e^{iks'} \Big((\mathsf{M}_1 + i\mathsf{N}_1) e^{-i\tilde{\beta}} + (\mathsf{M}_2 + i\mathsf{N}_2) e^{-i\tilde{\tilde{\beta}}} \Big) \,, \tag{3.9}$$

where $\tilde{\beta} = (k^2 \tilde{l}^{\,2})/(2\pi\tilde{\alpha})$ and $\tilde{\tilde{\beta}} = (k^2 \tilde{\tilde{l}}^{\,2})/(2\pi\tilde{\tilde{\alpha}})$; the values of $\tilde{X} \sim \tilde{Z}, \tilde{\tilde{X}} \sim \tilde{\tilde{Z}}$ are given from $X \sim Z$ using (3.7). [m]

The momentum field $\nabla \mathcal{S}_0$ is obtained from (3.9) as (there are implicit fixings of parameters in the following formulas; see p.24)

$$\mathbf{p} = \nabla \mathcal{S}_0 = \frac{\hbar}{i} \frac{\bar{\Psi}_{\text{diff}} \nabla \Psi_{\text{diff}} - \Psi_{\text{diff}} \nabla \bar{\Psi}_{\text{diff}}}{2\Psi_{\text{diff}} \Psi_{\text{diff}}} \,, \tag{3.10}$$

---

[m] $Ae^{-i\pi/4}/\sqrt{2(1-l^2)} \simeq Ae^{-i\pi/4}/\sqrt{2} = const.$ is omitted.
Note $\tilde{s}\,' = \sqrt{\tilde{X}^2 + \tilde{Y}^2 + \tilde{Z}^2} = \sqrt{\tilde{\tilde{X}}^2 + \tilde{\tilde{Y}}^2 + \tilde{\tilde{Z}}^2} = \tilde{\tilde{s}}\,' = s' = \sqrt{X^2 + Y^2 + Z^2}$.

We give explicit form of p_x ($p_y \simeq 0, p_z \simeq \hbar k_z$ is obvious). In

$$p_x = \frac{\partial \mathcal{S}_0}{\partial x} = \frac{\hbar}{i} \frac{\overline{\Psi}_{\mathrm{diff}} \partial_x \Psi_{\mathrm{diff}} - \Psi_{\mathrm{diff}} \partial_x \overline{\Psi}_{\mathrm{diff}}}{2 \Psi_{\mathrm{diff}} \overline{\Psi}_{\mathrm{diff}}} \ . \tag{3.11}$$

We have

$$\Psi_{\mathrm{diff}} \overline{\Psi}_{\mathrm{diff}} = \mathsf{M}_1^2 + \mathsf{N}_1^2 + \mathsf{M}_2^2 + \mathsf{N}_2^2 + 2(\mathsf{M}_1\mathsf{M}_2 + \mathsf{N}_1\mathsf{N}_2)\cos 2k_x x + 2(\mathsf{M}_1\mathsf{N}_2 - \mathsf{N}_1\mathsf{M}_2)\sin 2k_x x, \tag{3.12}$$

in which $\tilde{\beta} - \tilde{\tilde{\beta}} = -2k_x x$ was used, and

$$\begin{aligned} &\frac{\overline{\Psi}_{\mathrm{diff}} \partial_x \Psi_{\mathrm{diff}} - \Psi_{\mathrm{diff}} \partial_x \overline{\Psi}_{\mathrm{diff}}}{2i} = \mathrm{Im}(\overline{\Psi}_{\mathrm{diff}} \partial_x \Psi_{\mathrm{diff}}) \\ &= k_x(\mathsf{M}_1^2 + \mathsf{N}_1^2) - k_x(\mathsf{M}_2^2 + \mathsf{N}_2^2) + \mathsf{M}_1 \partial_x \mathsf{N}_1 - \mathsf{N}_1 \partial_x \mathsf{M}_1 + \mathsf{M}_2 \partial_x \mathsf{N}_2 - \mathsf{N}_2 \partial_x \mathsf{M}_2 \\ &\quad + \Big(- \mathsf{N}_1 \partial_x \mathsf{M}_2 + \mathsf{M}_1 \partial_x \mathsf{N}_2 - \mathsf{N}_2 \partial_x \mathsf{M}_1 + \mathsf{M}_2 \partial_x \mathsf{N}_1 \Big) \cos 2k_x x \\ &\quad + \Big(- \mathsf{M}_1 \partial_x \mathsf{M}_2 - \mathsf{N}_1 \partial_x \mathsf{N}_2 + \mathsf{M}_2 \partial_x \mathsf{M}_1 + \mathsf{N}_2 \partial_x \mathsf{N}_1 \Big) \sin 2k_x x \ , \end{aligned} \tag{3.13}$$

in which $\frac{\partial}{\partial x}(ks' - \tilde{\beta}) \simeq k_x$, $\frac{\partial}{\partial x}(ks' - \tilde{\tilde{\beta}}) \simeq -k_x$ was used. Thus we see that $p_x = \hbar \cdot$ RHS of (3.13)/RHS of (3.12). The forms of $\mathsf{M}_1, \mathsf{N}_1, \mathsf{M}_2$, and N_2 at $P = (X, 0, Z)$ on the coordinate system (x, y, z) are found in Appendix I.

• *Velocity field* The $\nabla \mathcal{S}_0$ is connected to $(\dot{\mathbf{q}}, \ddot{\mathbf{q}}, \dddot{\mathbf{q}})$ with (2.36) which is reduced to $\nabla \mathcal{S}_0 = m\dot{\mathbf{q}}$ since $\dot{\mathbf{q}}$ is large. Thus, the velocity field, which gives the particle's velocity at $\mathbf{x}$ *if* the particle comes to $\mathbf{x}$, is given as

$$\dot{x} = \frac{1}{m} \frac{\partial \mathcal{S}_0}{\partial x} = \frac{\hbar}{m} \frac{\text{RHS of (3.13)}}{\text{RHS of (3.12)}} \ , \quad \dot{y} = 0 \ , \quad \dot{z} = \frac{\hbar k_z}{m} \ . \tag{3.14}$$

• *Calculation of the electron distribution on the screen* We will calculate the electron distribution on the screen chasing trajectories obtained solving (3.14).

We assume that the incoming field is the two components Gaussian beam Ψ_s (see the sentences below (3.6)):

$$\Psi_s = \big(0.8 \exp(-\frac{(x-x_0)^2 + (y-y_0)^2}{0.34^{\ 2}}) + 0.2 \exp(-\frac{(x-x_0)^2 + (y-y_0)^2}{2.2^{\ 2}})\big) \times \exp(ikz) \ . \tag{3.15}$$

where the unit of length in (x, y)-plane is μm.

Since the Fresnel approximation breaks down in the vicinity of the biprism, we cannot follow the trajectory beyond the biprism. Therefore, 1) we have to assume some initial electron density distribution somewhere (our choice is $z = 4mm$) in the Fresnel region, and 2) we have to determine the diffracted field with which the electron located at (x_{ini}, y_{ini}) at $z = 4mm$ interacts, or equivalently, the center of the incoming beam from which the diffracted field originates. Regarding the 1), we assume that the electron density at $z = 4mm$ is equal to the field intensity formed by the uniform incoming Gaussian beam flux, which turns out to be equal to that formed by the incoming plane wave up to a const. factor. [n] Regarding the 2), we assume that the electron of which position is (x_{ini}, y_{ini}) at $z = 4mm$ interacts with the field which originates the Gaussian beam came along $(x_0 = x_{ini} - \frac{k_x}{k_z} \times 4mm,\ y_0 = y_{ini})$.

The electron density $I_{\text{lower}}(x)$ on the screen caused by lower-slot-passed electrons is reconstructed from the trajectories as follows. Let the initial x-coordinates of n sufficiently many electrons be $x_{1\cdot ini} < \cdots < x_{i\cdot ini} < \cdots < x_{n\cdot ini}$. The corresponding hitting x-coordinates be $x_{1\cdot hit} < \cdots < x_{i\cdot hit} < \cdots < x_{n\cdot hit}$. Let the field intensity at $z = 4mm$ be $|\Psi_{\text{diff}}(x)|^2$. Then I_{lower} at $x_{i\cdot hit}$ is given as

$$I_{\text{lower}}(x_{i\cdot hit}) \propto \frac{|\Psi_{\text{diff}}(x_{i\cdot ini})|^2}{\dot{x}|_{x_{i\cdot hit}}} \frac{(x_{(i+1)\cdot ini} - x_{(i-1)\cdot ini})}{(x_{(i+1)\cdot hit} - x_{(i-1)\cdot hit})} \quad (2 \le i \le n-1)\ . \tag{3.16}$$

The $I_{\text{upper}}(x)$ by upper-slot-passed electrons is given as $I_{\text{upper}}(x) = I_{\text{lower}}(-x)$.

Fig.3 is the assumed electron density at $z = 4mm$. Fig.5 shows the calculated trajectories of electrons.[o] The upper graph of Fig.4 is the electron density pattern calculated from the trajectories. [p] The middle graph is the field intensity calculated with the uniform incoming flux of two components Gaussian beam (3.15).[q] The lower graph is the field intensity calculated from the plane wave incoming beam. [r] Note that the envelope shapes are almost same in the middle and lower graphs. Note that the largest Fresnel peak of the electron density is shifted to the outerward at the arrowed positions. This is explained as follows. The electron runs almost linearly parallel to the geometrical border, whereas the Fresnel peak of the field runs along $\tilde{x} = const.\sqrt{\tilde{z}}$; see, e.g., Ref. 21 §8.7.3, see Fig. 5.

• *Velocity in the x-direction behind the biprism* As is seen from (3.14), the velocity in the x-direction in the i.f. region is slower in the both-slots-open case than that in the one-slot-open case ($k_x(\mathsf{M}_1^2 + \mathsf{N}_1^2) - k_x(\mathsf{M}_2^2 + \mathsf{N}_2^2)$ in the numerator is essential) because the oscillation energy is supplied from the x-component of the translational energy (see (3.1)); note 3-D QSHJE separates into three 1-D eqs. behind the biprism where the electric potential of the biprism disappears.

---

[n] The density of the lower-slot-passed electrons at $z = 4mm$ is $\mathsf{M}_1^2 + \mathsf{N}_1^2$ from (3.12). The M_1 and N_1 are given in (I.2). Note

$$\begin{aligned}\int_{-\infty}^{\infty} \Big(\int e^{-(\tau - x_0)^2} f(\tau) d\tau\Big)^2 dx_0 &= \int_{-\infty}^{\infty} dx_0 \int dy \int dz\, e^{-(y-x_0)^2} f(y) e^{-(z-x_0)^2} f(z) \\ &= \int f(y) dy \int f(z) dz \int_{-\infty}^{\infty} e^{-(y-x_0)^2} e^{-(z-x_0)^2} dx_0 = const.\Big(\int f(\tau) d\tau\Big)^2\ .\end{aligned}$$

[o] It took about 20 min. to calculate one trajectory on a 3GHz class personal computer with Mathematica.

[p] The upper graph was reconstructed from the trajectories of the lower-slot-passed electrons of which initial x-coordinates x_{ini}'s range from $x_{ini} = -2.000\mu m$ to $-0.105\mu m$, with $0.01\mu m$ step for $x_{ini} = -2.0\mu m \sim -0.930\mu m$ and with $0.005\mu m$ step for $x_{ini} = -0.925\mu m \sim -0.105\mu m$. The upper slot data were obtained reversing the lower slot data.

[q] The field intensity $I(x; x_0)$ caused by the two components Gaussian beam of which center (x, y) is $(x_0, 0)$ is obtained from (3.12) replacing $\mathsf{M}_1 \sim \mathsf{N}_2$ with $0.8\mathsf{M}_1 + 0.2\mathsf{M}_{11} \sim 0.8\mathsf{N}_2 + 0.2\mathsf{N}_{21}$, where what $\mathsf{M}_{11} \cdots \mathsf{N}_{21}$ represent are obvious. The field intensity $I(x)$ is obtained as $I(x) = \int_{-c}^{c} I(x; x_0) dx_0$, where c should be set sufficiently large. In the present case, we have set $c = 6.4\mu m$.

[r] The lower graph was calculated using (3.12) setting w_0 appearing in $\mathsf{M}_1 \sim \mathsf{N}_2$ to ∞.

We estimate the mean velocity in the x-direction at the screen position from the fringe visibility (f.v.) of the interference pattern. The f.v. is defined as f.v. $= (I_{max} - I_{min})/(I_{max} + I_{min})$, in which I_{max} (I_{min}) is the maximum (minimum) electron density on the screen. We consider that $I \propto |1/\dot{x}| \simeq m|\partial \mathcal{S}_0/\partial x|^{-1}$. For simplicity we assume that, in the i.f. region, the Ψ_{diff} is expressed as

$$\Psi_{\text{diff}} = e^{ikz}(J_{\text{up}}e^{ik_x x} + J_{\text{dn}}e^{-ik_x x}) , \tag{3.17}$$

in which $J_{\text{up}}, J_{\text{dn}} = const. > 0$. Then, $\partial \mathcal{S}_0/\partial x$ is given as

$$\frac{\partial \mathcal{S}_0}{\partial x} = \frac{\hbar k_x (J_{\text{up}}^2 - J_{\text{dn}}^2)}{J_{\text{up}}^2 + J_{\text{dn}}^2 + 2J_{\text{up}}J_{\text{dn}}\cos 2k_x x} . \tag{3.18}$$

From which we have f.v. $= 2J_{\text{up}}J_{\text{dn}}/(J_{\text{up}}^2 + J_{\text{dn}}^2)$. We define from (3.18) the mean velocity in the x-direction $\dot{x}_{mn}$ at the screen position as $m\dot{x}_{mn} = \hbar k_x (J_{\text{up}}^2 - J_{\text{dn}}^2)/(J_{\text{up}}^2 + J_{\text{dn}}^2)$. Thus, we have

$$\text{f.v.} = \sqrt{1 - (\frac{\dot{x}_{mn}}{\dot{x}_0})^2} , \tag{3.19}$$

where $\dot{x}_0 = \hbar k_x/m$. Some values are, for $\dot{x}_{mn}/\dot{x}_0 = 0.999$, f.v. $= 0.045$; for $\dot{x}_{mn}/\dot{x}_0 = 0.990$, f.v. $= 0.14$; for $\dot{x}_{mn}/\dot{x}_0 = 0.900$, f.v. $= 0.44$. The f.v. calculated from the upper graph of Fig.4 is $\approx 25\%$. Accordingly, the mean velocity in the x-direction at the screen position is, from (3.19), $\approx 97\%$ of that in the one-slot-open case. Thus, in the present case, the narrowing (compared to the one-slot-open case) of the envelope width of the i.f. pattern is not seen clearly.

• *Implicit assumption on* $A \sim D$ We calculated the trajectory assuming that an incoming beam accompanies a particle at its center. We however have no reason, at present, to exclude the possibility that the particle is located off-centered. In such a case, it can happen that the velocity field on the trajectory given with (3.14) changes sign at some point in the i.f. region. In other words, it can happen that the particle changes the moving direction from upward (downward) to downward (upward), which is what happens in B.I. (see Ref. 2 §5.1). In our formulation this moving direction change is not allowed because the J-O momentum $p_{(1)}$ is conserved in the region where no classical potential exists; see (C.6).

We show how we avoid the moving direction change. Remind, in 1-D space, the momentum $p_x = \partial \mathcal{S}_0/\partial x$ was determined, giving two linearly independent solutions $\Psi \in \mathbb{R}$ and $\Psi^D \in \mathbb{R}$ of the corresponding SE *and* Möbius parameters $A \sim D$, as (see (2.8))

$$p_x = \frac{\partial \mathcal{S}_0}{\partial x} = \frac{i}{2}\frac{\hbar(AD - BC)W}{(A\Psi^D + B\Psi)(C\Psi^D + D\Psi)} . \tag{3.20}$$

Also, we saw that the $\partial \mathcal{S}_0/\partial x$ determined from $(a + ic)\Psi^D + (b + id)\Psi = Re^{i\mathcal{S}_0/\hbar}$ is equal to that determined from (3.20) with $A = a + ic, B = b + id, C = a - ic, D = b - id$ (see p.7). We assume again for simplicity that the Ψ_{diff} in the i.f. region is given with (3.17). Then, writing x-component of Ψ_{diff} as $(J_{\text{up}} + J_{\text{down}})\cos k_x x + i(J_{\text{up}} - J_{\text{down}})\sin k_x x$, and identifying $(J_{\text{up}} + J_{\text{down}})\cos k_x x$ and $(J_{\text{up}} - J_{\text{down}})\sin k_x x$ as Ψ and Ψ^D of (3.20) respectively, we see that calculating the x-component p_x of the momentum using (3.18) is equivalent to choosing $\{A, B, C, D\} = \{i, 1, -i, 1\}$ in (3.20). If we choose $\{A, \cdots\} = \{-i, 1, i, 1\}$, the sign of p_x

is reversed ($p_x \to -p_x$). We can thus avoid the velocity reversal for the lower-slot-passed electron choosing $\{A, \cdots\} = \{i, 1, -i, 1\}$ if $J_{\rm up} > J_{\rm down}$, or $\{A, \cdots\} = \{-i, 1, i, 1\}$ if $J_{\rm up} < J_{\rm down}$. The choices for the upper-slot-passed electron are obvious.

Accordingly, the formula (3.14) is modified to

$$\dot{x} = \frac{1}{m}\frac{\partial \mathcal{S}_0}{\partial x} = \pm\frac{\hbar}{m}\,|\frac{\text{RHS of (3.13)}}{\text{RHS of (3.12)}}| \ , \quad \dot{y} = 0 \ , \quad \dot{z} = \frac{\hbar k_z}{m} \ , \tag{3.21}$$

where the '+ (-)' sign is for the particle which passed the lower (upper) slot. Note that the intervals on 1-D space (x-direction) divided by the point at which $p_x = 0$ are disconnected; the p_x is never zero in our formulation as is seen from (2.9). The values of the global consts. $A \sim D$ therefore need not be the same in each interval.

• *Experimental verification* We saw above that our formulation makes different predictions from QM. [s] Our formulation therefore testable. Examples of such predictions are
i) When the f.v. is small, the x-coordinate x_F on the screen of the largest Fresnel peak of the lower- (upper-) slot-passed electrons will increase (decrease) almost linearly as the distance z_{sc} between the screen and the biprism increases. While QM predicts that x_F increases (decreases) non-linearly as z_{sc} increases; see the paragraph below (3.16), see Fig.5.
ii) The larger the f.v. is, the narrower the overall i.f. pattern width is, which follows from (3.19). While QM predicts the overall i.f. pattern width is irrelevant to the value of the f.v.

We have encouraging data. On Fig.6, although the narrowing is not seen according to the small f.v. (≈ 0.02), the peak shifts seem to appear at the upper left and upper right corners. Similar peak shifts are seen also in Ref. 8 and Ref. 9.

---

[s] Even if the electron density in the (ensemble of the identical) incoming beam Ψ_s is assumed to be proportional to $|\Psi_s|^2$, the predictions below are still valid because the velocity in the x-direction is given not with (3.14) but with (3.21). If the electron density at $z = 4mm$ is given with $|\Psi_{\rm diff}|^2$ and the velocity is given with (3.14), the i.f. pattern (electron density) on the screen will be the same as the field intensity. Note if $m\dot{\mathbf{q}} = \nabla \mathcal{S}_0$, from the second eq. of (2.4), $R^2 = |\Psi_{\rm diff}|^2$ represents electron density.

4. Difference between Bohm's Interpretation and the Present Formulation

We first examine how our formulation differs from B.I., then show that the argument against introducing the trajectory to a relativistic spinless massive particle in B.I. does not apply to our formulation (introducing the trajectory to a relativistic Dirac particle in B.I. is possible; see, e.g., Ref. 2).

B.I. extracts the particle from the solution ψ of the time-independent SE of QM writing $\psi = Re^{i\mathcal{S}_0/\hbar}$ and interpreting $\nabla\mathcal{S}_0/m$ as the velocity of the particle: $\nabla\mathcal{S}_0 = m\dot{\mathbf{q}}$. While our formulation, as a modified CM, has the concept of the particle from the outset. In our formulation, the trajectory is determined with the E-L eq. (2.33) or with the eq. (2.36).

In B.I., the physical meaning of ψ in $\psi = Re^{i\mathcal{S}_0/\hbar}$ is the same as that given in QM since B.I. is an interpretation of QM. Accordingly, R^2 represents the particle density (in an ensemble consisting of systems of the same ψ). To make the particle number be time-independent in a time-independent potential, therefore, it has to be $\nabla\mathcal{S}_0 = m\dot{\mathbf{q}}$ according to the second eq. of (2.4). While in our formulation the physical meaning of $\psi = Re^{i\mathcal{S}_0/\hbar}$ is carried by the R and the $\mathcal{S}_0$ on the RHS. The $\mathcal{S}_0$ is a kind of 'field' defined all over $\mathbf{x}$ regardless of the particle position. If a particle comes to $\mathbf{x}$, $\nabla\mathcal{S}_0|_{\mathbf{x}}$ and $(\dot{\mathbf{q}}, \ddot{\mathbf{q}}, \dddot{\mathbf{q}})|_{\mathbf{x}}$ satisfy (2.36) there. The R also is regarded as a kind of 'field' in our formulation (see §3.3). At least, there is no reason for the R^2 to be equal to the particle density.

In B.I., the solution set $(R, \mathcal{S}_0)$ of (2.4) is acceptable if ψ in $\psi = Re^{i\mathcal{S}_0/\hbar}$ is quantum mechanically acceptable. The superposition of the different energy ψ's is therefore acceptable since it is acceptable in QM. While in our formulation, it is unacceptable. In our formulation, in accordance with CM, to a single action 'field' corresponds a single particle. In the case of a particle moving in a time-independent potential, the energy of the system is const. The superposition of different energy ψ's is therefore unphysical. Accordingly, the wave packet does not appear in our formulation, nor is it necessary.

In B.I., once the ψ is given, the $\nabla\mathcal{S}_0$ is uniquely determined from $\psi = Re^{i\mathcal{S}_0/\hbar}$. In our formulation, even if ψ is given, $\nabla\mathcal{S}_0$ is not uniquely determined. In 1-D space, $\nabla\mathcal{S}_0$ is determined giving ψ and the Möbius parameters $A \sim D$; see (2.8). For example, given $\psi = e^{ikx}$, B.I. gives uniquely $\nabla\mathcal{S}_0 = \hbar k$. While in our formulation, the $\nabla\mathcal{S}_0$ can be even $-\hbar k$ depending on the values of $A \sim D$; see (3.21).

In B.I., the velocity of the particle in a 1-D confining potential is zero (we restrict our discussion to 1-D space for simplicity). In B.I., the solution ψ of the 1-D SE of the system has to be an L^2-function since it has to be so in QM. Let $\Psi, \Psi^D \in \mathbb{R}$ be two linearly independent solutions of the SE. Let Ψ be the L^2-function. Then, Ψ^D is not an L^2-function (see i on p.6). Thus, in the form: $\psi = \Psi = Re^{i\mathcal{S}_0/\hbar}$, the $\mathcal{S}_0$ is const. since Ψ is real. From which $\dot{q} = \nabla\mathcal{S}_0/m \equiv 0$ follows (e.g., Ref. 2 §6.5.1). While in our formulation, it has to be $\psi = a\Psi + b\Psi^D$ with $0 \neq a, b \in \mathbb{C}$ (see §2.1). Accordingly, in the form: $\psi = Re^{i\mathcal{S}_0/\hbar}$, the $\mathcal{S}_0$ is not const. [3] From this $\mathcal{S}_0$, the velocity is obtained with the eq. (2.36).

Lastly, we show that the argument against the relativistic trajectory does not apply to our formulation. In the spirit of B.I., the relativistic trajectory of a spinless massive particle would be introduced as follows. i) Rewrite the unknown solution of the Klein-Gordon eq. (K-G eq.) as $\Phi(t, \mathbf{x}) = R(t, \mathbf{x})e^{iS(t,\mathbf{x})/\hbar}$. ii) Insert it into the K-G eq. iii) The resulting eq. is (2.3). iv) The four-velocity u^μ is given as $Mu^\mu = -(\partial^\mu S + (e/c)A^\mu)$, where $M = m(c^2 + Q)^{1/2}$, Q is given in (2.3a). The j^0 in (2.3b) is understood as the particle density in an ensemble. The reasons why this method does not work are i) j^0 is not positive definite, and ii) the velocity vector can be space-like; A space-like velocity is constructed using a superposition of the different energy solutions (e.g., Ref. 2 §12.1). In our formulation, however, i) j^0 does not represent the particle density, ii) in the time-independent electromagnetic field, the superposition of different energy solutions is unphysical (see the second last paragraph). Thus, it seems that we need not be so pessimistic about the possibility of the relativistic trajectory of the particle, although the theory construction is left to the future.

5. Concluding Remarks

We have formulated a quantum theory of a spinless massive particle as a generalized (or modified) analytical mechanics. i) We modified the classical H-J eq. to what we call the QSHJE using the extended action of the diffeomorphism group. ii) We determined the semiclassical Lagrangian consistent with the QSHJE; we found that the Lagrangian is of the second-order: $L(\mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}})$ and that, within our theoretical framework, there is no Lagrangian beyond the semiclassical level. iii) Based on the QSHJE and the Lagrangian, we explained the appearances of quantum phenomena relevant to the particle keeping the particle picture throughout.

Thus, one of the conceptual difficulties of quantum mechanics: the impossibility to maintain the particle picture, was resolved.

Our formulation is testable (with a present-day technology) because the particle density in an ensemble differs from that of quantum mechanics.

Acknowledgments

The author would like to express special gratitude to

A.E.Faraggi and M.Matone: If I had not happened to find their article, the present work did not even started. I appreciate their seminal works.

Toshio Ishigaki, ex. Professor of Hokkaido University: For guiding me through the world of the interpretations of Quantum Mechanics. For the warm support throughout the present work.

Kenzo Ishikawa, Professor of Hokkaido University: For providing me a research environment after T. Ishigaki's retirement. For guidance and discussions in the manuscript preparation stage, with which the present manuscript remarkably improved.



Fig. 3. Assumed electron density at $z = 4mm$.



Fig. 4. Interference patterns on the screen; Upper graph: electron density reconstructed from the trajectories; Middle graph: field intensity caused by the two components Gaussian beam (3.15); Lower graph: field intensity caused by the plane wave beam; Note that the electron density is larger than that expected in QM (that is, larger than the field intensity) at the arrowed positions. The marks on vertical axes are arbitrary.



Fig. 5. Trajectories of electrons which passed the lower slot. Crossed dotted lines borders on the geometrical shadows of the filament. The dotted curve is the path along which the largest peak of Fig.3 runs.



Fig. 6. Experimentally obtained electron density pattern and the calculated field intensity pattern reproduced from Ref. 7.

Appendix A. Extended action of $\mathrm{Diff}(\mathbb{RP}^1)$ on $\mathcal{F}_2(\mathbb{RP}^1)$

We give the mathematical background of the extended diffeomorphism action appeared in §2.1. In the following, we naively apply the results of the theory of finite-dimensional group extensions to the infinite-dimensional Lie group Diff($\mathbb{RP}^1$). For details on finite-dimensional group extensions, see Ref. 11, 12, 13, etc.

Let the diffeomorphism group of 1-D projective space $\mathbb{RP}^1$ be Diff($\mathbb{RP}^1$), the covariant tensor fields of order two (equivalently, the group of tensor densities of degree two) [t] on $\mathbb{RP}^1$ be $\mathcal{F}_2(\mathbb{RP}^1)$.

We consider the classical action T of $\mathrm{Diff}(\mathbb{RP}^1)$ on $\mathcal{F}_2(\mathbb{RP}^1)$ as a special case of more general action, in which T is given as $T : \mathrm{Diff}(\mathbb{RP}^1) \times \mathcal{F}_2(\mathcal{M}^1) \to \mathcal{F}_2(\mathbb{RP}^1),\ (y^{-1}, \phi(y)) \mapsto T_{y^{-1}}\phi(y) = \phi(y(x))(dy/dx)^2$. Let E be an extension of $\mathrm{Diff}(\mathbb{RP}^1)$ by $\mathcal{F}_2(\mathbb{RP}^1)$ or $0 \to \mathcal{F}_2(\mathbb{RP}^1) \to E \to \mathrm{Diff}(\mathbb{RP}^1) \to 1$ be a short exact sequence. Using the set-theoretic bijection between E and $\mathcal{F}_2(\mathbb{RP}^1) \times \mathrm{Diff}(\mathbb{RP}^1)$, we write an element of E as (ϕ, g), where $\phi \in \mathcal{F}_2(\mathbb{RP}^1)$ and $g \in \mathrm{Diff}(\mathbb{RP}^1)$. Let $s(g) := (dg, g) \in E$ and $i(\phi) := (\phi, 1) \in E$, where $s(g)$ is a section (a representative element of g in E) and $i(\phi)$ is an injection from $\mathcal{F}_2(\mathbb{RP}^1)$ to E. For a given action T, the operation rule in E is given [u] as $(\phi_1, g)(\phi_2, h) = (\phi_1 + T_g\phi_2 + f(g,h), gh)$, where $f(g,h)$ is a factor set which satisfies $s(g)s(h) = i(f(g,h))s(gh)$. As a diffeomorphism, we require the $s(g)$ to satisfy $s(g)s(h) = s(gh)$ or the above sequence to split; accordingly, $f(g,h) \equiv 0$. In E, the action of $\mathrm{Diff}(\mathbb{RP}^1)$ on $\mathcal{F}_2(\mathbb{RP}^1)$ is given as the product of $s(g)$ and $i(\phi)$: $s(g)i(\phi) = (dg, g)(\phi, 1) = (dg + T_g\phi, g)$, or $T_{(dg,g)}\phi = dg + T_g\phi$. If we set $dg = 0$, the classical action is recovered. In the present case that $\phi \in \mathcal{F}_2(\mathbb{RP}^1)$, we can set $dg^{-1} = \alpha\{g(x), x\}$, where $\alpha = const. \in \mathbb{R}$ and $\{g(x), x\}$ is the Schwarzian derivative, because it satisfies $d(g^{-1}h^{-1}) = dg^{-1} + T_{g^{-1}}dh^{-1}$ required from $s(g^{-1})s(h^{-1}) = s(g^{-1}h^{-1})$. In this case, the action $T_{(dg^{-1},\, g^{-1})}$ of $\mathrm{Diff}(\mathbb{RP}^1)$ on $\mathcal{F}_2(\mathbb{RP}^1)$ is given as [10] $T_{(dg^{-1},\, g^{-1})}\phi(y) = (dg/dx)^2\phi(g(x)) + \alpha\{g(x), x\}$.

---

[t] A covariant tensor field ϕ of order n on 1-D space $\mathcal{M}^1$ is written as $\phi = \phi(x)(dx)^n$, where x is a local coordinate. The field ϕ is also called a tensor density of degree n. Equivalently, a tensor density of degree n is defined as a section of the line bundle $(T^*\mathcal{M}^1)^{\otimes n}$, where $T^*\mathcal{M}^1$ is the cotangent bundle (e.g., Ref. 10).

[u] The group operation in $\mathcal{F}_n(\mathcal{M}^1)$ is written additively (e.g., $\phi_3 = \phi_1 + \phi_2$), that in $\mathrm{Diff}(\mathcal{M}^1)$ and E is written multiplicatively (e.g., $g_3 = g_1g_2$). Accordingly, the unit elements are written as 0, 1, and 1 respectively.

Appendix B. Derivation of the Relativistic Quantum H-J Equation

We derive (2.3). Our Minkowski metric is $\mathrm{diag}(1,-1,-1,-1)$. We use the Einstein summation convention. We work with the Gaussian system of electromagnetic units.

First, we examine the effect of the extended diffeomorphism action on the simplified relativistic H-J eq. We set A^{μ} in the relativistic H-J eq. of a charged particle: [22]

$$(\partial_{\mu}\mathcal{S}+\frac{e}{c}A_{\mu})(\partial^{\mu}\mathcal{S}+\frac{e}{c}A^{\mu})=\frac{1}{c^2}(\frac{\partial\mathcal{S}}{\partial t}+eA^0)^2-(\frac{\partial\mathcal{S}}{\partial\mathbf{x}}-\frac{e}{c}\mathbf{A})^2=m^2c^2 \tag{B.1}$$

to zeros, then restrict the eq. to 2-D time-space $(x^0(=ct),x^1)$ to obtain

$$\Big(\frac{1}{c}\frac{\partial\mathcal{S}(t,x^1)}{\partial t}\Big)^2-\Big(\frac{\partial\mathcal{S}(t,x^1)}{\partial x^1}\Big)^2=m^2c^2\ . \tag{B.2}$$

We examine the effect of the extended action of a diffeomorphism $y:x^1\mapsto y(x^1)$ upon the second term of the LHS of (B.2) with the first term left intact. Note, with the classical (not-extended) action, (B.2) is transformed to the eq. on the coordinate system (ct,y) as

$$\Big(\frac{1}{c}\frac{\partial\mathcal{S}(t,x^1(y))}{\partial t}\Big)^2-(\frac{\partial y}{\partial x^1})^2\Big(\frac{\partial\mathcal{S}(t,x^1(y))}{\partial y}\Big)^2=m^2c^2\ . \tag{B.3}$$

How is (B.2) transformed if the action is given with (2.1)? Let $q^a\to q^b\to q^c(=x^1)$ be a transformation among coordinate systems. By abuse of language, we use the same symbols $q^a,q^b,\cdots$ as labels of coordinate systems, as transformation functions, and as coordinate values. The coordinate transformation on 1-D space $\mathcal{M}^1$ from q^c to q^b is given by the action of $(dq^{c\ -1},q^{c\ -1})\in$ Ext. Diff$(\mathcal{M}^1)$ upon $(\partial\mathcal{S}/\partial q^c)^2\in\mathcal{F}_2(\mathcal{M}^1)$ as (see (2.1))

$$T_{(dq^{c\ -1},q^{c\ -1})}(\frac{\partial\mathcal{S}}{\partial q^c})^2=(\frac{\partial q^c}{\partial q^b})^2(\frac{\partial\mathcal{S}}{\partial q^c})^2+\alpha\{q^c,q^b\}=(\frac{\partial\mathcal{S}}{\partial q^b})^2+\alpha\{q^c,q^b\}, \tag{B.4}$$

where $\alpha=const.\in\mathbb{R}$. Successive transformation from q^b to q^a takes (B.4) to

$$\begin{aligned}T_{(dq^{b\ -1},q^{b\ -1})}T_{(dq^{c\ -1},q^{c\ -1})}(\frac{\partial\mathcal{S}}{\partial q^c})^2&=(\frac{\partial q^b}{\partial q^a})^2\Big((\frac{\partial\mathcal{S}}{\partial q^b})^2+\alpha\{q^c,q^b\}\Big)+\alpha\{q^b,q^a\}\\&=(\frac{\partial\mathcal{S}}{\partial q^a})^2+\alpha(\frac{\partial q^b}{\partial q^a})^2\{q^c,q^b\}+\alpha\{q^b,q^a\}\ .\end{aligned} \tag{B.5}$$

Using the cocycle condition: $\{q^c,q^a\}=(\partial q^b/\partial q^a)^2\{q^c,q^b\}+\{q^b,q^a\}$, we see that (B.5) is equal to

$$T_{(dq^{b\ -1},q^{b\ -1})}T_{(dq^{c\ -1},q^{c\ -1})}(\frac{\partial\mathcal{S}}{\partial q^c})^2=(\frac{\partial\mathcal{S}}{\partial q^a})^2+\alpha\{q^c,q^a\}\ . \tag{B.6}$$

Thus, for the coordinate transformation $q^a\to q^b\to q^c$, the eq. (B.2) on q^c:

$$(\frac{1}{c}\frac{\partial\mathcal{S}}{\partial t})^2-(\frac{\partial\mathcal{S}}{\partial q^c})^2=m^2c^2$$

is transformed to the eq. on q^b as

$$(\frac{1}{c}\frac{\partial\mathcal{S}}{\partial t})^2-(\frac{\partial q^b}{\partial q^c})^2\Big((\frac{\partial\mathcal{S}}{\partial q^b})^2+\alpha\{q^c,q^b\}\Big)=m^2c^2\ . \tag{B.7}$$

Then, (B.7) is transformed to the eq. on q^a as

$$(\frac{1}{c}\frac{\partial \mathcal{S}}{\partial t})^2-(\frac{\partial q^a}{\partial q^c})^2\Big((\frac{\partial \mathcal{S}}{\partial q^a})^2+\alpha\{q^c,q^a\}\Big)=m^2c^2\ . \tag{B.8}$$

While the eq. on q^c is considered as the transformed eq. from that on $q^\star$. Thus, the eqs. on q^c, q^b, and q^a take the same form:

$$(\frac{1}{c}\frac{\partial \mathcal{S}}{\partial t})^2-(\frac{\partial q^c}{\partial q^c})^2\Big((\frac{\partial \mathcal{S}}{\partial q^c})^2+\alpha\{q^\star,q^c\}\Big)=m^2c^2 \tag{B.9a}$$

$$(\frac{1}{c}\frac{\partial \mathcal{S}}{\partial t})^2-(\frac{\partial q^b}{\partial q^c})^2\Big((\frac{\partial \mathcal{S}}{\partial q^b})^2+\alpha\{q^\star,q^b\}\Big)=m^2c^2 \tag{B.9b}$$

$$(\frac{1}{c}\frac{\partial \mathcal{S}}{\partial t})^2-(\frac{\partial q^a}{\partial q^c})^2\Big((\frac{\partial \mathcal{S}}{\partial q^a})^2+\alpha\{q^\star,q^a\}\Big)=m^2c^2\ . \tag{B.9c}$$

However, an equation upon the coordinate system $q^c(q^b,q^a)$ cannot contain a transformation function $q^\star$ to another coordinate $q^\star$. The $q^\star$ therefore has to be replaced with some quantity defined on $q^c(q^b,q^a)$. In (B.9a), if $q^\star=Aq^c+B$ (for which $\{q^\star,q^c\}=0$), $\mathcal{S}=-cp_0t+p_1q^c$, where $p_0=const.>0$ and $p_1=const.\neq 0$, is a solution, which hints at replacing $q^\star$ with $\mathcal{S}$. This replacement is supported by the following statement: A physical quantity defined on 2-D Minkowski time-space (ct,q^c) which is relevant to a free particle and linear in q^c is almost singled out as $\mathcal{S}$. Accordingly, we set $q^\star=\mathcal{S}$. [v] Thus, if the action is given with (2.1), (B.2) is transformed to

$$(\frac{1}{c}\frac{\partial \mathcal{S}}{\partial t})^2-\Big((\frac{\partial \mathcal{S}}{\partial x^1})^2+\alpha\{\mathcal{S},x^1\}\Big)=m^2c^2\ . \tag{B.10}$$

When an electromagnetic field exists, (B.9a) becomes

$$(\frac{1}{c}\frac{\partial \mathcal{S}}{\partial t}+\frac{e}{c}A^0)^2-\Big((\frac{\partial \mathcal{S}}{\partial q^c}-\frac{e}{c}A^1)^2+\alpha\{q^\star,q^c\}\Big)=m^2c^2\ . \tag{B.11}$$

With the same reasoning as that in the last paragraph, $\partial_1 q^\star$ is singled out as $\partial_1\mathcal{S}-\frac{e}{c}A^1$. We therefore replace $\partial_1 q^\star$ with$\partial_1\mathcal{S}-\frac{e}{c}A^1$. Thus, (B.11) is written as

$$m^2c^2=(\frac{1}{c}\frac{\partial \mathcal{S}}{\partial t}+\frac{e}{c}A^0)^2-\Big((\frac{\partial \mathcal{S}}{\partial x^1}-\frac{e}{c}A^1)^2+\alpha\big(\frac{\partial_1^2(\partial_1\mathcal{S}-\frac{e}{c}A^1)}{(\partial_1\mathcal{S}-\frac{e}{c}A^1)}-\frac{3}{2}\big(\frac{\partial_1(\partial_1\mathcal{S}-\frac{e}{c}A^1)}{(\partial_1\mathcal{S}-\frac{e}{c}A^1)}\big)^2\big)\Big)\ . \tag{B.12}$$

Next work is to determine the 4-D relativistic Quantum H-J eq.

First, we will find the eq. valid for the electromagnetic potential A's which take the form: $A=(A^0(x^0),A^1(x^1),A^2(x^2),A^3(x^3))$. Let a set of all electromagnetic potential A's which take such forms on a Lorentz frame x be $\mathcal{A}_{\text{sep}}$. If $A\in\mathcal{A}_{\text{sep}}$, (B.1) separates into four 1-D eqs. with $\mathcal{S}=\mathcal{S}_0(x^0)+\mathcal{S}_1(x^1)+\mathcal{S}_2(x^2)+\mathcal{S}_3(x^3)$ on the frame. [w] On the assumption that the extended diffeomorphism group acts upon $\mathcal{F}_2(\mathcal{M}^1)$, each 1-D eq. is modified to the form similar to (B.12). Collecting each modified eqs., we have

---

[v] Because $q^\star\in\text{Diff}(\mathcal{M}^1)$, as a result of equating $\mathcal{S}$ with $q^\star$, $\mathcal{S}$ (seen as a function of $q^c(q^b,q^a)$ at any fixed time t) is required to be, at least, monotonic.

[w] The idea to use $A=(A^0(x^0),\cdots,A^3(x^3))$ was borrowed from Ref. 23, in which Bertoldi et al. derived the 4-D relativistic quantum H-J eq. from ‘the Equivalence Principle’ elaborately explained in Ref. 3.

$$\Big((\frac{\partial \mathcal{S}}{\partial x^0} + \frac{e}{c} A^0)^2 + \alpha_0 Q^0 \Big) - \Big(\sum_{i=1}^{3} (\frac{\partial \mathcal{S}}{\partial x^i} - \frac{e}{c} A^i)^2 + \alpha_1 Q^1 + \alpha_2 Q^2 + \alpha_3 Q^3 \Big) = m^2 c^2 \ , \tag{B.13}$$

where $\alpha_0, \cdots, \alpha_3 \in \mathbb{R}$ and

$$Q^0 = \frac{\partial_0^2 (\partial_0 \mathcal{S} + \frac{e}{c} A^0)}{(\partial_0 \mathcal{S} + \frac{e}{c} A^0)} - \frac{3}{2} \Big(\frac{\partial_0 (\partial_0 \mathcal{S} + \frac{e}{c} A^0)}{(\partial_0 \mathcal{S} + \frac{e}{c} A^0)} \Big)^2 \ , \quad Q^i = \frac{\partial_i^2 (\partial_i \mathcal{S} - \frac{e}{c} A^i)}{(\partial_i \mathcal{S} - \frac{e}{c} A^i)} - \frac{3}{2} \Big(\frac{\partial_i (\partial_i \mathcal{S} - \frac{e}{c} A^i)}{(\partial_i \mathcal{S} - \frac{e}{c} A^i)} \Big)^2$$

($i = 1, 2, 3$; take no sum). Interestingly, say, Q^1 (likewise for Q^0, Q^2, Q^3) is written as $Q^1 = 2\partial_1 \partial^1 R_1 / R_1$ with the solution R_1 of $\partial_1 (R_1^2 (\partial_1 \mathcal{S}_1 - eA^1/c)) = 0$. [x] Then, considering $\alpha_0 = \alpha_i = \hbar^2/2$ is realized in nature (the resulting eq. becomes Lorentz invariant if $\alpha_0 = \alpha_i$, reproduces experimental data if $\alpha_0 = \alpha_i = \hbar^2/2$), we sum up non-classical terms $Q := \alpha_0 Q^0 - \alpha_i Q^i$ in (B.13) to $-\hbar^2 \partial_\mu \partial^\mu R/R$, where $R = R_0(x^0) R_1(x^1) R_2(x^2) R_3(x^3)$. The (B.13) is therefore rewritten in the form:

$$(\partial_\mu \mathcal{S} + \frac{e}{c} A_\mu)(\partial^\mu \mathcal{S} + \frac{e}{c} A^\mu) = m^2 c^2 + \frac{\hbar^2 \partial_\mu \partial^\mu R}{R} \tag{B.14a}$$

$$\partial_\mu j^\mu = 0 \ , \qquad j^\mu = -\frac{R^2}{m} (\partial^\mu \mathcal{S} + \frac{e}{c} A^\mu) \ . \tag{B.14b}$$

The (B.14) is a system of equations of which unknown functions are $\mathcal{S}$ and R. Although (B.14) looks Lorentz invariant, the invariance is not assured for $A \notin \mathcal{A}_{\text{sep}}$ since (B.14) is nothing but a rewriting of (B.13).

Next, we will find the eq. set valid not only for $A \in \mathcal{A}_{\text{sep}}$ but for arbitrary A's, that is, for $A^\mu = A^\mu(x^0, x^1, x^2, x^3)$. We assume that such an eq. set takes the form:

$$(\partial_\mu \mathcal{S} + \frac{e}{c} A_\mu)(\partial^\mu \mathcal{S} + \frac{e}{c} A^\mu) = m^2 c^2 + \frac{\hbar^2 \partial_\mu \partial^\mu R}{R} + g \tag{B.15a}$$

$$\partial_\mu j^\mu = \breve{g} \ , \qquad j^\mu = -\frac{R^2}{m} (\partial^\mu \mathcal{S} + \frac{e}{c} A^\mu) \ , \tag{B.15b}$$

where $g(\breve{g})$ is a Lorentz scaler which vanish ($g(\breve{g}) \equiv 0$) at least when $A \in \mathcal{A}_{\text{sep}}$. The $g(\breve{g})$ depends on some of the Lorentz scalers constructed from $\partial_\mu \mathcal{S}$, A, R and their derivatives. The eqs. to be gauge invariant, the A appears in the form: $(\partial^\mu \mathcal{S} + \frac{e}{c} A^\mu)$. Thus, g (likewise for $\breve{g}$) looks like $g(x) = g(\partial_\mu (\partial^\mu \mathcal{S} + \frac{e}{c} A^\mu)(x), (\partial \mathcal{S} + \frac{e}{c} A)^2(x), R(x), \partial_\mu \partial^\mu R(x), \partial_\mu R \partial^\mu R(x), \cdots)$. Because $g(\breve{g}) \equiv 0$ for $A \in \mathcal{A}_{\text{sep}}$ (including $A = 0$), and the functional form of $g(\breve{g})$ is common to $A \in \mathcal{A}_{\text{sep}}$ and $A \notin \mathcal{A}_{\text{sep}}$, $g(\breve{g}) \equiv 0$ for arbitrary A.

Thus, we see (B.14) as the modified H-J eq. on the assumption that the extended diffeomorphism group acts on each term of the LHS of (B.1) and call (B.14) the relativistic Quantum H-J equation.

---

[x] From $\partial_1 \big(R_1^2 (\partial_1 \mathcal{S}_1 - \frac{e}{c} A^1) \big) = 0$, if $\partial_1 \mathcal{S}_1 - \frac{e}{c} A^1 \neq 0$, $R_1 = \pm const. \, (\partial_1 \mathcal{S}_1 - \frac{e}{c} A^1)^{-1/2}$. Therefore

$$\frac{\partial_1 \partial^1 R_1}{R_1} = -\frac{\partial_1 \partial_1 R_1}{R_1} = \frac{1}{2} \Big(\frac{\partial_1^2 (\partial_1 \mathcal{S}_1 - \frac{e}{c} A^1)}{\partial_1 \mathcal{S}_1 - \frac{e}{c} A^1} - \frac{3}{2} \big(\frac{\partial_1 (\partial_1 \mathcal{S}_1 - \frac{e}{c} A^1)}{\partial_1 \mathcal{S}_1 - \frac{e}{c} A^1} \big)^2 \Big) \ .$$

Appendix C. Higher-Order Analytical Mechanics

We summarize the 3rd order analytical mechanics, in which the Lagrangian is a function of $\mathbf{q}$, $\dot{\mathbf{q}}$, $\ddot{\mathbf{q}}$, and $\dddot{\mathbf{q}}$. Although it is possible to consider still higher-order analytical mechanics, we don't need them. Materials in this section are found in Ref. 5 and 6. References 24 and 25 are useful as supporting readings. See them for details.

• *Lagrangian formalism* ^{5} As in CM, action $\mathcal{S}$ is defined as $\mathcal{S} := \int_a^b L(\mathbf{q},\dot{\mathbf{q}},\ddot{\mathbf{q}},\dddot{\mathbf{q}})dt$. Let $\delta\mathbf{V}$ be the variation of the curve $\mathbf{q}(t)$. Then, we have

$$\begin{aligned}
\delta\mathcal{S} &= \int L(\mathbf{q}+\delta\mathbf{V},\dot{\mathbf{q}}+\delta\dot{\mathbf{V}},\ddot{\mathbf{q}}+\delta\,\ddot{\mathbf{V}}\,,\dddot{\mathbf{q}}+\delta\,\dddot{\mathbf{V}}\,)dt - \int L(\mathbf{q},\dot{\mathbf{q}},\ddot{\mathbf{q}},\dddot{\mathbf{q}})dt \\
&= \int (\frac{\partial L}{\partial \mathbf{q}}\delta\mathbf{V} + \frac{\partial L}{\partial \dot{\mathbf{q}}}\delta\dot{\mathbf{V}} + \frac{\partial L}{\partial \ddot{\mathbf{q}}}\delta\ddot{\mathbf{V}} + \frac{\partial L}{\partial \dddot{\mathbf{q}}}\delta\dddot{\mathbf{V}})dt = \int (\frac{\partial L}{\partial \mathbf{q}} - \frac{d}{dt}\frac{\partial L}{\partial \dot{\mathbf{q}}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \ddot{\mathbf{q}}} - \frac{d^3}{dt^3}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\delta\mathbf{V}dt \\
&\quad + (\frac{\partial L}{\partial \dot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \ddot{\mathbf{q}}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\delta\mathbf{V}|_a^b + (\frac{\partial L}{\partial \ddot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\delta\dot{\mathbf{V}}|_a^b + (\frac{\partial L}{\partial \dddot{\mathbf{q}}})\delta\ddot{\mathbf{V}}|_a^b \,. \qquad \text{(C.1)}
\end{aligned}$$

From the first term in the RHS of (C.1), we obtain the E-L equation:

$$\frac{\partial L}{\partial \mathbf{q}} - \frac{d}{dt}\frac{\partial L}{\partial \dot{\mathbf{q}}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \ddot{\mathbf{q}}} - \frac{d^3}{dt^3}\frac{\partial L}{\partial \dddot{\mathbf{q}}} = 0 \,. \qquad \text{(C.2)}$$

We define $\mathbf{p}_{(1)}$, $\mathbf{p}_{(2)}$, and $\mathbf{p}_{(3)}$ as (See (C.1))

$$\mathbf{p}_{(1)} := \frac{\partial L}{\partial \dot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \ddot{\mathbf{q}}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \dddot{\mathbf{q}}} \,, \qquad \mathbf{p}_{(2)} := \frac{\partial L}{\partial \ddot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \dddot{\mathbf{q}}} \,, \qquad \mathbf{p}_{(3)} := \frac{\partial L}{\partial \dddot{\mathbf{q}}} \,. \qquad \text{(C.3)}$$

The $\mathbf{p}_{(1)}$, $\mathbf{p}_{(2)}$, and $\mathbf{p}_{(3)}$ are called Jacobi-Ostrogradski (J-O) momenta. We define $\overset{3}{\mathcal{E}}_c\,(L)$ as

$$\begin{aligned}
\overset{3}{\mathcal{E}}_c\,(L) &= \mathbf{p}_{(1)}\dot{\mathbf{q}} + \mathbf{p}_{(2)}\ddot{\mathbf{q}} + \mathbf{p}_{(3)}\dddot{\mathbf{q}} - L \\
&= (\frac{\partial L}{\partial \dot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \ddot{\mathbf{q}}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\dot{\mathbf{q}} + (\frac{\partial L}{\partial \ddot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\ddot{\mathbf{q}} + \frac{\partial L}{\partial \dddot{\mathbf{q}}}\dddot{\mathbf{q}} - L \,. \qquad \text{(C.4)}
\end{aligned}$$

The $\overset{3}{\mathcal{E}}_c\,(L)$, which is reduced to $(\partial L/\partial\dot{\mathbf{q}})\dot{\mathbf{q}} - L$ if $L = L(\mathbf{q},\dot{\mathbf{q}})$, is const. along the solution curve of the E-L eq. (C.2) as is seen below:

$$\begin{aligned}
&\frac{d}{dt}\overset{3}{\mathcal{E}}_c\,(L) = \frac{d}{dt}(\mathbf{p}_{(1)}\dot{\mathbf{q}} + \mathbf{p}_{(2)}\ddot{\mathbf{q}} + \mathbf{p}_{(3)}\dddot{\mathbf{q}} - L) \\
&= (\frac{d}{dt}\frac{\partial L}{\partial \dot{\mathbf{q}}} - \frac{d^2}{dt^2}\frac{\partial L}{\partial \ddot{\mathbf{q}}} + \frac{d^3}{dt^3}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\dot{\mathbf{q}} + (\frac{\partial L}{\partial \dot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \ddot{\mathbf{q}}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\ddot{\mathbf{q}} \\
&+ (\frac{d}{dt}\frac{\partial L}{\partial \ddot{\mathbf{q}}} - \frac{d^2}{dt^2}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\ddot{\mathbf{q}} + (\frac{\partial L}{\partial \ddot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\dddot{\mathbf{q}} + (\frac{d}{dt}\frac{\partial L}{\partial \dddot{\mathbf{q}}})\dddot{\mathbf{q}} + (\frac{\partial L}{\partial \dddot{\mathbf{q}}})\ddddot{\mathbf{q}} \\
&- \dot{\mathbf{q}}\frac{\partial L}{\partial \mathbf{q}} - \ddot{\mathbf{q}}\frac{\partial L}{\partial \dot{\mathbf{q}}} - \dddot{\mathbf{q}}\frac{\partial L}{\partial \ddot{\mathbf{q}}} - \ddddot{\mathbf{q}}\frac{\partial L}{\partial \dddot{\mathbf{q}}} = -\dot{\mathbf{q}}\left(\frac{\partial L}{\partial \mathbf{q}} - \frac{d}{dt}\frac{\partial L}{\partial \dot{\mathbf{q}}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \ddot{\mathbf{q}}} - \frac{d^3}{dt^3}\frac{\partial L}{\partial \dddot{\mathbf{q}}}\right) = 0 \,. \qquad \text{(C.5)}
\end{aligned}$$

We use also the symbol E for $\overset{3}{\mathcal{E}}_c\,(L)$. Along the solution curve of (C.2), we have

$$\frac{d\mathbf{p}_{(1)}}{dt} - \frac{\partial L}{\partial \mathbf{q}} = \frac{d}{dt}(\frac{\partial L}{\partial \dot{\mathbf{q}}} - \frac{d}{dt}\frac{\partial L}{\partial \ddot{\mathbf{q}}} + \frac{d^2}{dt^2}\frac{\partial L}{\partial \dddot{\mathbf{q}}}) - \frac{\partial L}{\partial \mathbf{q}} = -\frac{\partial L}{\partial \mathbf{q}} + \frac{d}{dt}\frac{\partial L}{\partial \dot{\mathbf{q}}} - \frac{d^2}{dt^2}\frac{\partial L}{\partial \ddot{\mathbf{q}}} + \frac{d^3}{dt^3}\frac{\partial L}{\partial \dddot{\mathbf{q}}} = 0 \,. \qquad \text{(C.6)}$$

If the Lagrangian does not contain $\mathbf{q}$ explicitly, therefore, $d\mathbf{p}_{(1)}/dt = 0$.

- *Hamiltonian formalism* [6] We use a symbol $\overset{(m)}{\mathbf{q}}$ for $d^m\mathbf{q}/dt^m$ $(m = 0, 1, \cdots)$; $\overset{(0)}{\mathbf{q}} = \mathbf{q}$. If the Lagrangian is regular $(|\partial^2 L/\partial \dddot{q}^i \partial \dddot{q}^j| \neq 0)$, $Leg : T^5M \to T^*(T^2M), (\mathbf{q}, \overset{(1)}{\mathbf{q}}, \cdots, \overset{(5)}{\mathbf{q}}) \mapsto (\mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}}, \mathbf{p}_{(1)}, \mathbf{p}_{(2)}, \mathbf{p}_{(3)})$ is a local diffeomorphism called the Legendre transformation; where T^5M is the 5th order tangent bundle and $T^*(T^2M)$ is the cotangent bundle on T^2M. Note the 6th order E-L eq. obtained from the 3rd order regular Lagrangian determines a vector field on T^5M. Let $H = \overset{3}{\mathcal{E}}_c\,(L) \circ Leg^{-1}$. Comparing two expressions

$$dH = \frac{\partial H}{\partial \overset{(n-1)}{\mathbf{q}}} d \overset{(n-1)}{\mathbf{q}} + \frac{\partial H}{\partial \mathbf{p}_{(n)}} d\mathbf{p}_{(n)}\ , \quad dH = -\frac{\partial L}{\partial \mathbf{q}} d\mathbf{q} + \overset{(n)}{\mathbf{q}}\, d\mathbf{p}_{(n)} + \big(\mathbf{p}_{(n)} - \frac{\partial L}{\partial \overset{(n)}{\mathbf{q}}}\big) d \overset{(n)}{\mathbf{q}}$$

$(n = 1, 2, 3)$, we have, using (C.2) and (C.3),

$$\frac{d \overset{(n-1)}{\mathbf{q}}}{dt} = \frac{\partial H}{\partial \mathbf{p}_{(n)}}\ , \quad \frac{d\mathbf{p}_{(n)}}{dt} = -\frac{\partial H}{\partial \overset{(n-1)}{\mathbf{q}}} \quad (n = 1, 2, 3).$$

Appendix D. The c_k may depend on the integration parameters of the E-L eq.

The formulas (2.20), (2.21) and (2.22) take the same forms in both cases: i) c_k's $(k = 1, 2a, 2b, 3, \cdots)$ are consts. ii) c_k's depend on the start point $(x_0(t_0), \dot{x}_0(t_0))$ and the end point $(x(t_1), \dot{x}(t_1))$ of the action integral. Note the dependence on which is equivalent to the dependence on the initial conditions $(q(t_0), \cdots, \dddot{q}(t_0))$ of the 1-D E-L eq. or on the $\{v, a_1, b_1, q_0 (:= \text{initial position})\}$ of (2.26) since any of them is a set of four integration parameters of the 4th order E-L eq.

If c_k's depend on $(x_0(t_0), \dot{x}_0(t_0))$ and $(x(t_1), \dot{x}(t_1))$, the (C.1) becomes, for a 1-D Lagrangian $L(q, \dot{q}, \ddot{q}; c_k(x, \dot{x}))$ (we use a 2nd order Lagrangian for simplicity; we fix the start point $(x_0, \dot{x}_0)$),

$$\begin{aligned}
\delta\mathcal{S} &= \int^{x(t_1), \dot{x}(t_1)} L(q + \delta V, \dot{q} + \delta\dot{V}, \ddot{q} + \delta\ddot{V}; c_k(x, \dot{x}))dt - \int^{x(t_1), \dot{x}(t_1)} L(q, \dot{q}, \ddot{q}; c_k(x, \dot{x}))dt \\
&= \int^{x(t_1), \dot{x}(t_1)} ((\delta V \frac{\partial}{\partial q} + \delta\dot{V} \frac{\partial}{\partial \dot{q}} + \delta\ddot{V} \frac{\partial}{\partial \ddot{q}}) L(\cdots; c_k(x, \dot{x}))dt \\
&= \int^{x(t_1), \dot{x}(t_1)} (\frac{\partial}{\partial q} - \frac{d}{dt}\frac{\partial}{\partial \dot{q}} + \frac{d^2}{dt^2}\frac{\partial}{\partial \ddot{q}}) L(\cdots; c_k(x, \dot{x}))\delta V dt \\
&\quad + (\frac{\partial}{\partial \dot{q}} - \frac{d}{dt}\frac{\partial}{\partial \ddot{q}}) L(\cdots; c_k(x + \delta x, \dot{x}))\delta V|^{x(t_1), \dot{x}(t_1)} + \frac{\partial L(\cdots; c_k(x, \dot{x} + \delta\dot{x}))}{\partial \ddot{q}} \delta\dot{V}|^{x(t_1), \dot{x}(t_1)}\ ,
\end{aligned}$$

where $\delta x = \delta V|_{(x, \dot{x})}$, $\delta\dot{x} = \delta\dot{V}|_{(x, \dot{x})}$. Because the end point $(x, \dot{x})$ is fixed in the E-L eq. determination process, the resulting E-L eq. is the same in both cases. That $p_{(1)}$ has the same form in both cases is seen from

$$p_{(1)} = \lim_{\delta x \to 0} \frac{\delta\mathcal{S}}{\delta x} = \lim_{\delta x \to 0} (\frac{\partial}{\partial \dot{q}} - \frac{d}{dt}\frac{\partial}{\partial \ddot{q}}) L(\cdots; c_k(x + \delta x, \dot{x})) = (\frac{\partial}{\partial \dot{q}} - \frac{d}{dt}\frac{\partial}{\partial \ddot{q}}) L(\cdots; c_k(x, \dot{x}))\ .$$

Similarly for $p_{(2)}$.

Appendix E. Characteristics of the solution of the 1-D E-L Equation

We examine the solution of the 1-D E-L eq. (2.20).

• *The velocity never vanishes* The $\dot{q}$ of a solution of the E-L eq. (2.20) never take value zero unless $\ddot{q} \equiv 0$ as is shown as follows. Let $q(t)$ be a solution of the E-L eq. Then,

$$p_{(1)} = m\dot{q} + \sum_{k=2}^{+\infty} \frac{c_k \hbar^k}{m^{k-1}} \Big(- \frac{k(k-1)\dddot{q}\ddot{q}^{k-2}}{\dot{q}^{3k-2}} + \frac{(3k-2)(k-1)\ddot{q}^k}{\dot{q}^{3k-1}} \Big) \tag{E.1}$$

$$E = \frac{1}{2}m\dot{q}^2 + \sum_{k=2}^{+\infty} \frac{c_k \hbar^k}{m^{k-1}} \Big(- \frac{k(k-1)\dddot{q}\ddot{q}^{k-2}}{\dot{q}^{3k-3}} + \frac{(3k-1)(k-1)\ddot{q}^k}{\dot{q}^{3k-2}} \Big) , \tag{E.2}$$

where $c_2 = 3c_{2a} + c_{2b}$, are conserved (cf. (2.21a) and (2.22)). Removing $\dddot{q}$, we have

$$\frac{1}{2}m\dot{q}^2 - p_{(1)}\dot{q} + E = \sum_{k=2}^{+\infty} \frac{(k-1)c_k \hbar^k}{m^{k-1}} \frac{\ddot{q}^k}{\dot{q}^{3k-2}} \ . \tag{E.3}$$

Assume $\dot{q}(t) \to 0$ for $\ddot{q}(t) \neq 0$. For $\dot{q} \to 0$, no term of the RHS of (E.3) approaches $\pm\infty$; otherwise, the LHS cannot be kept finite because the convergence speed of each term of the RHS is different. Thus, $|\ddot{q}^k/\dot{q}^{3k-2}| \leq const.$ for $\dot{q} \to 0$. From which we see $\ddot{q}$ approaches zero faster than or equal to $\dot{q}^2$. Consequently the $\dot{q}$ never take value zero.

• *The velocity is written as a Fourier series* We show, for any semiclassical initial conditions (p.10), the solution of the E-L eq. (2.20) of which $c_2 = 3c_{2a} + c_{2b} < 0$ is written as $q(t) = vt + \sum_{n=1}^{\infty} A_n \cos(n\omega t + \theta_n)$. Since initial conditions are semiclassical, we have $|\ddot{q}(t_0)| << m|\dot{q}(t_0)|^3/(\hbar\sqrt{-2c_2}\,)$ and $|\dddot{q}(t_0)| << m^2|\dot{q}(t_0)|^5/(-2\hbar^2 c_2)$ by definition. Accordingly, all terms in the RHS of (2.20) except $m\ddot{q}$ and $-2c_2\hbar^2\ddddot{q}/(m\dot{q}^4)$ are small compared to the $m\ddot{q}$. The (2.20) is therefore written approximately as $0 = m\ddot{q} - 2c_2\hbar^2\ddddot{q}/(m\dot{q})$. From an approximate solution $q(t) = vt + \tilde{A}\cos(\tilde{\omega}t + \tilde{\theta})$ of it, we see that $\ddot{q}$ takes value zero repeatedly and at time t at which $\ddot{q} = 0$, $\dddot{q} \neq 0$ (otherwise $q(t) = vt$ is the solution). Inserting $\ddot{q} = 0$ into (E.1) and (E.2), and solving resulting eqs. for $\dot{q}$ and $\dddot{q}$, we have $\dot{q} = (p_{(1)} \pm \sqrt{p_{(1)}^2 - 2mE}\,)/m$ and $\dddot{q} = \pm(p_{(1)} \pm \sqrt{p_{(1)}^2 - 2mE}\,)^4 \sqrt{p_{(1)}^2 - 2mE}/(2c_2\hbar^2 m^3)$. Since $p_{(1)}$ and E are *consts*., at times at which $\dot{q} = (p_{(1)} + \sqrt{p_{(1)}^2 - 2mE}\,)/m$, the values of $\dot{q}$, $\ddot{q}(=0)$, and $\dddot{q}$ are equal respectively, which means the oscillation is periodic and the $\tilde{\omega}$ is the fundamental angular frequency (freq.). Thus, the $\dot{q}(t)$ is written as a Fourier series.

• *The inf. order of A_n is $(\dot{f}/v)^n$* Since the $\tilde{\omega}$ is the fundamental angular freq., $\tilde{A}\cos(\tilde{\omega}t + \tilde{\theta}) = A_1\cos(\omega t + \theta_1)$ and $\omega A_2/v, \omega A_3/v, \cdots \in \mathcal{O}(> \dot{f}/v)$. The lowest inf. order oscillating term in $(\dot{f}/v)^2$-terms: $2c_2\hbar^2 \cdot 8\ddot{q}\dddot{q}/(m\dot{q}^5)$ and $-2c_3\hbar^3(3\dddot{q}^2 + 3\ddot{q}\ddddot{q})/(m^2\dot{q}^7)$ of (2.20) is $const. \cdot A_1^2\cos(2\omega t + 2\theta_1 + \xi)$ generated from $A_1^2\omega^5\cos(\omega t + \theta_1)\sin(\omega t + \theta_1)$, $A_1^2\omega^6\sin^2(\omega t + \theta_1)$ and $A_1^2\omega^6\cos^2(\omega t + \theta_1)$. It sums up to zero with the 2ω term: $const. \cdot A_2\cos(2\omega t + \theta_2)$ coming from $m\ddot{q}$ and $-2c_2\hbar^2\ddddot{q}/m\dot{q}^4$. Thus, we see $A_2 \in \mathcal{O}((\dot{f}/v)^2)$. In a similar manner, we see $A_n \in \mathcal{O}((\dot{f}/v)^n)$.

Appendix F. Determination of c_{2b} in the 1-D Semiclassical Lagrangian

We determine c_{2b} in the 1-D Lagrangian (2.19) of which $c_1 = c_{2a} = 0$. We also determine v, a_1, b_1 and ω of the trajectory (2.26) as functions of ε_1, ε_2, and k of p_{QHJ} (2.28). They are determined as follows. From the E-L eq., we obtain $\omega = mv^2/(\hbar\sqrt{-2c_{2b}})$. It gives $c_{2b} = -mv^2/(4E)$ because $\omega = kv$ (see next paragraph). The v in $c_{2b} = -mv^2/(4E)$ is determined as a function of a_1, b_1 and E from (2.22). Lastly the a_1 and b_1 are determined from the comparison of p_{QHJ} and p_{EL}.

• *Determination of ω* We determine the relation between $k = \sqrt{2mE}/\hbar$ and ω comparing p_{QHJ} (2.28) and p_{EL} (2.23). We expand p_{QHJ} of (2.28) as

$$\begin{aligned} p_{QHJ} &= \sqrt{2mE}\big(1 - \varepsilon_1 \cos 2kx - \varepsilon_2 \sin 2kx + \mathcal{O}(\varepsilon_1^2) + \mathcal{O}(\varepsilon_2^2) + \mathcal{O}(\varepsilon_1\varepsilon_2)\big) \\ &= \sqrt{2mE}\big(1 - \varepsilon_1 \cos 2kx - \varepsilon_2 \sin 2kx + \mathcal{O}(\varepsilon^2)\big) \; . \end{aligned} \tag{F.1}$$

Because we are in the semiclassical world, $\varepsilon_1, \varepsilon_2 << 1$; we wrote the $\mathcal{O}(\varepsilon_1^2) + \mathcal{O}(\varepsilon_2^2) + \mathcal{O}(\varepsilon_1\varepsilon_2)$ as $\mathcal{O}(\varepsilon^2)$ with $\varepsilon = \sqrt{\varepsilon_1^2 + \varepsilon_2^2}$, which is allowed since ε_1^2, ε_2^2, $|\varepsilon_1\varepsilon_2| \le (\varepsilon_1^2 + \varepsilon_2^2)$. We expand p_{EL} of (2.23), inserting $c_1 = c_{2a} = 0$ and $q = vt + f$, as

$$p_{EL} = p_{(1)} + \frac{2c_{2b}\hbar^2}{m}\frac{\ddot{f}^2}{(v + \dot{f})^5} + \mathcal{O}(\hbar^3) \; . \tag{F.2}$$

We note that $\mathcal{O}(\hbar^3)$ of (F.2) is equal to $\mathcal{O}((\dot{f}/v)^3)$. We know, among the oscillating terms in (2.26), $A_1 \cos(\omega t + \theta_1)$ has the lowest inf. order with respect to (w.r.t.) $\dot{f}/v$ (Appendix E). The angular freq. of the oscillation of p_{EL} having the lowest inf. order w.r.t. $\dot{f}/v$ is therefore 2ω generated from $A_1^2 \cos^2(\omega t + \theta_1)$. While the wave number of the oscillation of p_{QHJ} having the lowest inf. order w.r.t. ε is $2k$. We therefore have $\omega = kv$. In passing, we see that $\mathcal{O}(\varepsilon) = \mathcal{O}((\dot{f}/v)^2)$.

• *Determination of c_{2b}* We insert $c_1 = c_{2a} = 0$ and $q(t)$ of (2.26) into the E-L eq. (2.20). Then, we have

$$-m\ddot{f} + \frac{2c_{2b}\hbar^2}{m}(\frac{\ddddot{f}}{v^4}) + \mathcal{O}(|\frac{\dot{f}}{v}|^2) = (m\omega^2 + \frac{2c_{2b}\hbar^2\omega^4}{mv^4})A_1 \cos(\omega t + \theta_1) + \mathcal{O}(|\frac{\dot{f}}{v}|^2) = 0 \; .$$

From which we have

$$c_{2b} = \frac{-m^2v^4}{2\hbar^2\omega^2} = \frac{-mv^2}{4E} \; . \tag{F.3}$$

• *Determination of $p_{(1)}$ and E as functions of a_1, b_1, ω and v* Using $\dot{f}\dot{f} = \omega^2((a_1^2 + b_1^2)/2 - a_1 b_1 \sin 2\omega t + (b_1^2 - a_1^2)\cos 2\omega t/2) + \mathcal{O}(\delta^3)$, and so forth, where $\delta := \dot{f}/v$, we extract non-oscillating terms from $p_{(1)}$ and E:

$$\begin{aligned} p_{(1)} &= m\dot{q} - \frac{2c_{2b}\hbar^2}{m}\frac{\dddot{q}}{\dot{q}^4} + \frac{4c_{2b}\hbar^2}{m}\frac{\ddot{q}^2}{\dot{q}^5} + \mathcal{O}(\hbar^3) = m(v + \dot{f}) \\ &\quad - \frac{2c_{2b}\hbar^2}{m}(\frac{\dddot{f}}{v^4} - \frac{4\dddot{f}\dot{f}}{v^5} + \frac{10\dddot{f}\dot{f}^2}{v^6} + \cdots) + \frac{4c_{2b}\hbar^2}{m}(\frac{\ddot{f}^2}{v^5} - \frac{5\ddot{f}^2\dot{f}}{v^6} + \cdots) + \mathcal{O}(\hbar^3) \end{aligned} \tag{F.4}$$

$$E=\frac{1}{2}m\dot{q}^2-\frac{2c_{2b}\hbar^2}{m}\frac{\dddot{q}}{\dot{q}^3}+\frac{5c_{2b}\hbar^2}{m}\frac{\ddot{q}^2}{\dot{q}^4}+\mathcal{O}(\hbar^3)=\frac{1}{2}m(v^2+2v\dot{f}+\dot{f}\dot{f})$$
$$-\frac{2c_{2b}\hbar^2}{m}\Big(\frac{\dddot{f}}{v^3}-\frac{3\dddot{f}\dot{f}}{v^4}+\frac{6\dddot{f}\dot{f}^2}{v^5}+\cdots\Big)+\frac{5c_{2b}\hbar^2}{m}\Big(\frac{\ddot{f}^2}{v^4}-\frac{4\ddot{f}^2\dot{f}}{v^5}+\cdots\Big)+\mathcal{O}(\hbar^3) \qquad \text{(F.5)}$$

at the inf. orders up to δ^3 to have (the $\hbar^3$- and $\hbar^4$-terms of (2.21a) and (2.22) do not generate non-oscillating terms)

$$mv-\frac{2c_{2b}\hbar^2}{m}\Big(\frac{-4}{v^5}\Big)\frac{(-\omega^4)(a_1^2+b_1^2)}{2}+\frac{4c_{2b}\hbar^2}{mv^5}\frac{\omega^4(a_1^2+b_1^2)}{2}+\mathcal{O}(\delta^4)$$
$$=mv+\frac{4\hbar^2\omega^4}{mv^5}\frac{a_1^2+b_1^2}{2}(-c_{2b})+\mathcal{O}(\delta^4)=p_{(1)}=const. \qquad \text{(F.6)}$$

$$\frac{1}{2}mv^2+\frac{1}{2}m\frac{\omega^2(a_1^2+b_1^2)}{2}-\frac{2c_{2b}\hbar^2}{m}\Big(\frac{-3}{v^4}\Big)\frac{(-\omega^4)(a_1^2+b_1^2)}{2}+\frac{5c_{2b}\hbar^2}{mv^4}\frac{\omega^4(a_1^2+b_1^2)}{2}+\mathcal{O}(\delta^4)$$
$$=\frac{m}{2}v^2+\frac{m\omega^2}{2}\frac{(a_1^2+b_1^2)}{2}+\frac{\hbar^2\omega^4}{mv^4}\frac{(a_1^2+b_1^2)}{2}(-c_{2b})+\mathcal{O}(\delta^4)=E=const. \qquad \text{(F.7)}$$

- *The a_1,b_1 as functions of ε_1, ε_2, ω and v* To express a_1,b_1 of (F.6) and (F.7) as functions of ε_1, ε_2 and E ($a_1=a_1(\varepsilon_1,\varepsilon_2,E), b_1=b_1(\varepsilon_1,\varepsilon_2,E)$), we once again compare p_{QHJ} and p_{EL}. We compare (F.2) written in the form:

$$p_{EL}=p_{(1)}+\frac{2c_{2b}\hbar^2}{mv^5}\Big(\omega^2\sqrt{a_1^2+b_1^2}\cos(\omega t+\theta_1)\Big)^2+\mathcal{O}(\delta^3)$$
$$=p_{(1)}+\frac{2c_{2b}\hbar^2\omega^4}{mv^5}\Big(\frac{a_1^2+b_1^2}{2}+a_1b_1\sin 2\omega t+\frac{a_1^2-b_1^2}{2}\cos 2\omega t\Big)+\mathcal{O}(\delta^3) \qquad \text{(F.8)}$$

with (F.1) written in the form

$$p_{QHJ}=\sqrt{2mE}\big(1-\varepsilon_1\cos 2kx-\varepsilon_2\sin 2kx+\mathcal{O}(\varepsilon^2)\big)$$
$$=\sqrt{2mE}\big(1-\varepsilon_1\cos 2k(vt+f)-\varepsilon_2\sin 2k(vt+f)+\mathcal{O}(\varepsilon^2)\big)=\sqrt{2mE}\times$$
$$\big(1-\varepsilon_1\cos 2\omega t-\varepsilon_2\sin 2\omega t-2kf(\varepsilon_2\cos 2\omega t-\varepsilon_1\sin 2\omega t)+\mathcal{O}((kf)^2\varepsilon)+\mathcal{O}(\varepsilon^2)\big)\ . \qquad \text{(F.9)}$$

to have

$$\frac{2c_{2b}\hbar^2\omega^4}{mv^5}a_1b_1+\mathcal{O}(\delta^4)=-\sqrt{2mE}\ (\varepsilon_2+\mathcal{O}(\varepsilon^2)) \qquad \text{(F.10a)}$$
$$\frac{2c_{2b}\hbar^2\omega^4}{mv^5}\frac{a_1^2-b_1^2}{2}+\mathcal{O}(\delta^4)=-\sqrt{2mE}\ (\varepsilon_1+\mathcal{O}(\varepsilon^2))\ . \qquad \text{(F.10b)}$$

Note $kf\simeq\dot{f}/v<<1$, which is seen as follows. The oscillation amplitude f is roughly given as increasing rate $\dot{f}$ times half-cycle time $\pi/(2\omega)$: $f\simeq\dot{f}\times\pi/(2\omega)$. Using $\omega=kv$, we see $2kf/\pi\simeq\dot{f}/v$. Note there is no term having angular freq. 2ω at the inf. order δ^3 in p_{QHJ} of (F.9) nor in p_{EL} of (F.8). We therefore converted $\mathcal{O}(\delta^3)$ in (F.8) to $\mathcal{O}(\delta^4)$ in (F.10). From (F.10) we have (double sign corresponds)

$$a_1^2=\frac{mv^5\sqrt{2mE}}{2c_{2b}\hbar^2\omega^4}\big(-\varepsilon_1\pm\sqrt{\varepsilon_1^2+\varepsilon_2^2}\ +\mathcal{O}(\varepsilon^2)\big),\ b_1^2=\frac{mv^5\sqrt{2mE}}{2c_{2b}\hbar^2\omega^4}\big(\varepsilon_1\pm\sqrt{\varepsilon_1^2+\varepsilon_2^2}\ +\mathcal{O}(\varepsilon^2)\big).$$

- *Summary up to this point* Thus, we have obtained (remind $\varepsilon = \sqrt{\varepsilon_1^2 + \varepsilon_2^2}$)

$$\omega = kv \ , \quad c_{2b} = \frac{-m^2 v^4}{2\hbar^2 \omega^2} = \frac{-mv^2}{4E} \tag{F.11a}$$

$$a_1^2 = \frac{mv^5 \sqrt{2mE}}{2c_{2b}\hbar^2\omega^4}(-\varepsilon_1 \pm \varepsilon + \mathcal{O}(\varepsilon^2)) \ , \quad b_1^2 = \frac{mv^5 \sqrt{2mE}}{2c_{2b}\hbar^2\omega^4}(\varepsilon_1 \pm \varepsilon + \mathcal{O}(\varepsilon^2)) \tag{F.11b}$$

$$p_{(1)} = mv + \frac{4\hbar^2\omega^4}{mv^5}\frac{a_1^2 + b_1^2}{2}(-c_{2b}) + \mathcal{O}(\delta^4) \tag{F.11c}$$

$$E = \frac{m}{2}v^2 + \frac{m\omega^2}{2}\frac{(a_1^2 + b_1^2)}{2} + \frac{\hbar^2\omega^4}{mv^4}\frac{(a_1^2 + b_1^2)}{2}(-c_{2b}) + \mathcal{O}(\delta^4) \ . \tag{F.11d}$$

- *Expressing c_{2b}, v, ω, a_1, b_1, and $p_{(1)}$ as functions of $\varepsilon_1, \varepsilon_2$, and E* We express each term in (F.11) as a function of $\varepsilon_1, \varepsilon_2$, and E. We assume $v > 0$. Then the double sign in (F.11b) is fixed to minus. Thus,

$$a_1^2 = \frac{\hbar^2}{mv\sqrt{2mE}}(\varepsilon_1 + \varepsilon + \mathcal{O}(\varepsilon^2)), \quad b_1^2 = \frac{\hbar^2}{mv\sqrt{2mE}}(-\varepsilon_1 + \varepsilon + \mathcal{O}(\varepsilon^2)) \ . \tag{F.12}$$

Inserting (F.12) into (F.11d), and solving for v, that is solving $mv^2/2 + \sqrt{2mE}(\varepsilon + \mathcal{O}(\varepsilon^2))v - E = 0$, we have

$$v = \frac{\sqrt{2mE}}{m}\big(\sqrt{1 + (\varepsilon + \mathcal{O}(\varepsilon^2))^2} - (\varepsilon + \mathcal{O}(\varepsilon^2))\big) = \frac{\sqrt{2mE}}{m}(1 - \varepsilon + \mathcal{O}(\varepsilon^2)) \ . \tag{F.13}$$

From (F.13), we have

$$c_{2b} = \frac{-mv^2}{4E} = -\frac{1}{2}(1 - 2\varepsilon + \mathcal{O}(\varepsilon^2)) \ , \quad \omega = kv = \frac{2E}{\hbar}(1 - \varepsilon + \mathcal{O}(\varepsilon^2)) \tag{F.14}$$

$$p_{(1)} = \sqrt{2mE}(1 + \varepsilon + \mathcal{O}(\varepsilon^2)) \ . \tag{F.15}$$

Inserting v of (F.13) into a_1^2, b_1^2 of (F.12), we have

$$a_1^2 = -\frac{\hbar^2}{2mE}(1 + \varepsilon)(-\varepsilon_1 - \varepsilon) = -\frac{\hbar^2}{2mE}(-\varepsilon_1 - \varepsilon + \mathcal{O}(\varepsilon^2))$$
$$b_1^2 = -\frac{\hbar^2}{2mE}(1 + \varepsilon)(\varepsilon_1 - \varepsilon) = -\frac{\hbar^2}{2mE}(\varepsilon_1 - \varepsilon + \mathcal{O}(\varepsilon^2)) \ .$$

However, it has to be (see (F.10a))

$$-\varepsilon_2\sqrt{2mE} = \frac{2c_{2b}\hbar^2\omega^4}{mv^5}a_1 b_1 = -\frac{\hbar^2}{2mE}(1 + \varepsilon)a_1 b_1 \ .$$

Therefore, if $\varepsilon_2 \geq 0$ (double sign corresponds)

$$a_1 = \pm\sqrt{-\frac{\hbar^2}{2mE}(-\varepsilon_1 - \varepsilon)} + \mathcal{O}(\varepsilon^{\frac{3}{2}}), \ b_1 = \pm\sqrt{-\frac{\hbar^2}{2mE}(\varepsilon_1 - \varepsilon)} + \mathcal{O}(\varepsilon^{\frac{3}{2}}). \tag{F.16}$$

If $\varepsilon_2 < 0$ (double sign corresponds)

$$a_1 = \pm\sqrt{-\frac{\hbar^2}{2mE}(-\varepsilon_1 - \varepsilon)} + \mathcal{O}(\varepsilon^{\frac{3}{2}}), \ b_1 = \mp\sqrt{-\frac{\hbar^2}{2mE}(\varepsilon_1 - \varepsilon)} + \mathcal{O}(\varepsilon^{\frac{3}{2}}). \tag{F.17}$$

Appendix G. Inconsistency between p_{QHJ} and p_{EL} at $(\dot{f}/v)^3$

To see the inconsistency, we compare the p_{QHJ} and p_{EL} at $\varepsilon^{3/2} = (\dot{f}/v)^3$. We have

$$p_{QHJ} = \frac{\hbar k(1+\varepsilon_1)}{(\sin kx + \varepsilon_2 \cos kx)^2 + (1+\varepsilon_1)^2 \cos^2 kx} = \hbar k\big(1 - \varepsilon(1 - \frac{\varepsilon_1}{2})\cos(2kx - \theta_Q)\big)$$

$$+\mathcal{O}(\varepsilon^2) = \hbar k\big(1 - \varepsilon(1 - \frac{\varepsilon_1}{2})\cos(2kvt + 2kf - \theta_Q)\big) + \mathcal{O}(\varepsilon^2)$$

$$= \hbar k\big(1 - \varepsilon(1 - \frac{\varepsilon_1}{2})\cos(2\omega t - \theta_Q) - 2\varepsilon k f \sin(2\omega t - \theta_Q)\big) + \mathcal{O}(\varepsilon^2)\ ,$$

where the θ_Q satisfies $\cos\theta_Q = (\varepsilon^2/2 + \varepsilon_1)/\sqrt{(\varepsilon^2/2+\varepsilon_1)^2 + \varepsilon_2^2}$ and $\sin\theta_Q = \varepsilon_2/\sqrt{(\varepsilon^2/2+\varepsilon_1)^2 + \varepsilon_2^2}$; remind $\varepsilon = \sqrt{\varepsilon_1^2 + \varepsilon_2^2}$. While we have

$$p_{EL} = p_{(1)} + \frac{2c_{2b}\hbar^2}{m}(\frac{\ddot{f}^2}{v^5} - \frac{5\dot{f}\ddot{f}^2}{v^6}) + \frac{3c_3\hbar^3}{m^2}\frac{\ddot{f}^3}{v^8} + \mathcal{O}((\frac{\dot{f}}{v})^4)\ .$$

The 3ω-term at $\varepsilon^{3/2}$ of p_{QHJ} is

$$-\hbar k\varepsilon k A_1 \sin(3\omega t + \theta_1 - \theta_Q) = -\frac{mEk^3A_1^3}{\hbar k}\sin(3\omega t + 3\theta_1)\ , \tag{G.1}$$

where we used $\varepsilon = mEA_1^2/\hbar^2$ ((F.16) or (F.17)) and $\theta_Q = -2\theta_1$ obtained from the comparison of p_{QHJ} and p_{EL} at $\varepsilon^1 = (\dot{f}/v)^2$. While the 3ω-terms at $(\dot{f}/v)^3$ of p_{EL} is

$$\begin{aligned}&\frac{2c_{2b}\hbar^2\omega^2}{mv^3}\Big(\frac{4\omega^2A_1A_2}{v^2}\cos(3\omega t + \theta_1 + \theta_2) + \frac{5\omega^3A_1^3}{4v^3}\sin(3\omega t + 3\theta_1)\Big)\\ &\qquad - \frac{3c_3\hbar^3\omega^3}{4m^2v^5}\frac{\omega^3A_1^3}{v^3}\cos(3\omega t + 3\theta_1) = -\frac{\hbar^2k^2}{mv}kA_1\times\\ &\Big(4kA_2\cos(3\omega t + \theta_1 + \theta_2) + \frac{5}{4}k^2A_1^2\sin(3\omega t + 3\theta_1) + \frac{3c_3\hbar k}{4mv}k^2A_1^2\cos(3\omega t + 3\theta_1)\Big)\ ,\end{aligned} \tag{G.2}$$

where we used $\omega = kv$, $2\theta_1 = -\theta_Q$ and $c_{2b} = -1/2 + \mathcal{O}(\varepsilon)$. Inserting the E-L eq.'s solution (2.27) with c_2 in which replaced with c_{2b}, we rewrite (G.2) as

$$(\text{G.2}) \Rightarrow -\frac{\hbar^2k^2}{mv}k^3A_1^3\big((\frac{-2}{(-2c_{2b})^{\frac{3}{2}}} + \frac{3\hbar k}{4mv})c_3\cos(3\omega t + 3\theta_1) + \frac{13}{4}\sin(3\omega t + 3\theta_1)\big). \tag{G.3}$$

We cannot make both of the amplitude and the phase of (G.3) to be equal to those of (G.1) adjusting c_3. In a similar manner, we can show that ω-terms at $(\dot{f}/v)^3$ of p_{EL} cannot be made equal to ω-terms at $\varepsilon^{3/2}$ of p_{QHJ}, though we don't work it out here.

Appendix H. Determination of c_{2b}^{3D} of the 3-D free Lagrangian

We determine c_{2b}^{3D} of the 3-D Lagrangian $L = m\dot{\mathbf{q}}^2/2 + c_{2b}^{3D}\hbar^2\ddot{\mathbf{q}}^2/m\dot{\mathbf{q}}^4 + c_4^{3D}\hbar^4\ddot{\mathbf{q}}^4/m^3\dot{\mathbf{q}}^{10} + \mathcal{O}(\hbar^5)$. We set $c_3^{3D} = 0$ (p.12).

First, we prepare expressions of the E-L eq., $\mathbf{p}_{EL}$ and E obtained from the Lagrangian above (see (C.2), (2.36) and (C.4)):

$$m\ddot{q}_i + \frac{c_{2b}^{3D}\hbar^2}{m}\Big(-\frac{2\ddddot{q}_i}{\dot{\mathbf{q}}^4} + \frac{16\dot{\mathbf{q}}\ddot{\mathbf{q}}\dddot{q}_i}{\dot{\mathbf{q}}^6} - \frac{8\ddot{\mathbf{q}}\dddot{\mathbf{q}}\dot{q}_i}{\dot{\mathbf{q}}^6} + \frac{24\ddot{\mathbf{q}}^2(\dot{\mathbf{q}}\ddot{\mathbf{q}})\dot{q}_i}{\dot{\mathbf{q}}^8} + \frac{(4\ddot{\mathbf{q}}^2 + 8\dot{\mathbf{q}}\dddot{\mathbf{q}})\ddot{q}_i}{\dot{\mathbf{q}}^6} - \frac{48(\dot{\mathbf{q}}\ddot{\mathbf{q}})^2\ddot{q}_i}{\dot{\mathbf{q}}^8}\Big) + \mathcal{O}(\hbar^4) = 0 \tag{H.1}$$

$$p_{EL\,i} = p_{(1)i} + p_{(2)i}\frac{\ddot{q}_i}{\dot{q}_i} = p_{(1)i} + \frac{2c_{2b}^{3D}\hbar^2\ddot{q}_i^2}{m\dot{\mathbf{q}}^4\dot{q}_i} + \frac{4c_4^{3D}\hbar^4\ddot{\mathbf{q}}^2\ddot{q}_i^2}{m^3\dot{\mathbf{q}}^{10}\dot{q}_i} + \cdots \tag{H.2}$$

$$E = \mathbf{p}_{(1)}\dot{\mathbf{q}} + \mathbf{p}_{(2)}\ddot{\mathbf{q}} - L = \frac{1}{2}m\dot{\mathbf{q}}^2 + \frac{\hbar^2 c_{2b}^{3D}}{m}\Big(-\frac{2\dot{\mathbf{q}}\dddot{\mathbf{q}}}{\dot{\mathbf{q}}^4} - \frac{3\ddot{\mathbf{q}}^2}{\dot{\mathbf{q}}^4} + \frac{8(\dot{\mathbf{q}}\ddot{\mathbf{q}})^2}{\dot{\mathbf{q}}^6}\Big) + \frac{\hbar^4 c_4^{3D}}{m^3}\Big(-\frac{7\ddot{\mathbf{q}}^4}{\dot{\mathbf{q}}^{10}} - \frac{8(\dot{\mathbf{q}}\ddot{\mathbf{q}})(\ddot{\mathbf{q}}\dddot{\mathbf{q}})}{\dot{\mathbf{q}}^{10}} - \frac{4\ddot{\mathbf{q}}^2(\dot{\mathbf{q}}\dddot{\mathbf{q}})}{\dot{\mathbf{q}}^{10}} + \frac{40\ddot{\mathbf{q}}^2(\dot{\mathbf{q}}\ddot{\mathbf{q}})^2}{\dot{\mathbf{q}}^{12}}\Big) + \cdots\ , \tag{H.3}$$

We determine c_{2b}^{3D}. First, we see $c_{2b}^{3D} < 0$; otherwise a wavy linear trajectory cannot be a solution. For the negative c_{2b}^{3D} and a semiclassical initial conditions, the solution of (H.1) is written as $q_i(t) = v_i t + A_{i1}\cos(\omega t + \theta_{i1}) + g_i(t)\ \cdots(*1)$, where $v_i = const. > 0, g_i \in \mathcal{O}(\boldsymbol{\delta}^2)$, $\boldsymbol{\delta} := \omega|\ \mathbf{A}_1|/|\mathbf{v}| << 1, |\ \mathbf{A}_1| = \sqrt{(}A_{11}^2 + A_{21}^2 + A_{31}^2)$, likewise for $|\mathbf{v}|$. From the truncated E-L eq.: $m\ddot{q}_i - 2\hbar^2 c_{2b}^{3D}\ddddot{q}_i/(m\dot{\mathbf{q}}^4) = 0\ \cdots(*2)$, we have $\omega = m\mathbf{v}^2/(\hbar\sqrt{(}-2c_{2b}^{3D}))$. Note $g_i(t) \neq \sum_{n=2}^{+\infty} A_{in}\cos(n\omega t + \theta_{in})$ in general. Indeed, if we insert $q_i(t) = v_i t + \sum_{n=1}^{+\infty} A_{in}\cos(n\omega t + \theta_{in})$ into (H.1), non-oscillating terms appearing at the inf. order $\boldsymbol{\delta}^2$ does not vanish in general. Inserting $\omega = \mathbf{k}\mathbf{v} = \sum_{i=1}^{3} v_i\sqrt{2mE_i}/\hbar$ into the ω above, we have

$$c_{2b}^{3D} = -\frac{m^2\mathbf{v}^4}{2\hbar^2\omega^2} = \frac{-m^2(v_1^2 + v_2^2 + v_3^2)^2}{2(v_1\sqrt{2mE_1} + v_2\sqrt{2mE_2} + v_3\sqrt{2mE_3})^2}\ . \tag{H.4}$$

To express v_i as the function of ε_{1i}, ε_{2i}, and E_i of (2.31), first, we write A_{i1} as a function of $(\varepsilon_{1i}, \varepsilon_{2i})$. Then, we write v_i as a function of A_{i1} and E_i. We insert $(*1)$ into (H.2) to have

$$\begin{aligned} p_{EL\,i} &= p_{(1)i} + p_{(2)i}\frac{\ddot{q}_i}{\dot{q}_i} = const. + \frac{\hbar^2}{m}\frac{2c_{2b}^{3D}\ddot{q}_i^2}{\dot{\mathbf{q}}^4\dot{q}_i} + \mathcal{O}(\boldsymbol{\delta}^4) \\ &= const. + \frac{\hbar^2}{m}\frac{2c_{2b}^{3D}\omega^2}{\mathbf{v}^2 v_i}\Big(\frac{\omega^2}{\mathbf{v}^2}(\frac{a_{i1}^2 + b_{i1}^2}{2} + a_{i1}b_{i1}\sin 2\omega t + \frac{a_{i1}^2 - b_{i1}^2}{2}\cos 2\omega t) + \mathcal{O}(\boldsymbol{\delta}^3)\Big)\ , \end{aligned}$$

where $a_{i1} = A_{i1}\cos\theta_{i1}$ and $b_{i1} = -A_{i1}\sin\theta_{i1}$. Comparing it with $p_{QHJ\,i}$ (cf. (F.9)):

$$p_{QHJ\,i} = \sqrt{2mE_i}\big(1 - \varepsilon_{1i}\cos 2\omega t - \varepsilon_{2i}\sin 2\omega t + \mathcal{O}(\boldsymbol{\varepsilon}^{\frac{3}{2}})\big)$$

at the inf. order $\boldsymbol{\delta}^2 = \boldsymbol{\varepsilon}^1$, where $\boldsymbol{\varepsilon} = \sum_{i=1}^{3} \sqrt{(\varepsilon_{1i}^2 + \varepsilon_{2i}^2)}$, we have

$$\frac{\hbar^2}{m}\frac{2c_{2b}^{3D}\omega^2}{\mathbf{v}^2 v_i}\Big(\frac{\omega^2}{\mathbf{v}^2}\frac{(a_{i1}^2 - b_{i1}^2)}{2} + \mathcal{O}(\boldsymbol{\delta}^3)\Big) = -\varepsilon_{1i}\sqrt{2mE_i}\ ,$$
$$\frac{\hbar^2}{m}\frac{2c_{2b}^{3D}\omega^2}{\mathbf{v}^2 v_i}\Big(\frac{\omega^2}{\mathbf{v}^2}a_{i1}b_{i1} + \mathcal{O}(\boldsymbol{\delta}^3)\Big) = -\varepsilon_{2i}\sqrt{2mE_i}\ ,$$

from which we have

$$\frac{m\omega^2 A_{i1}^2}{2v_i\sqrt{2mE_i}} + \mathcal{O}(\boldsymbol{\delta}^3) = \frac{m\omega^2(a_{i1}^2 + b_{i1}^2)}{2v_i\sqrt{2mE_i}} + \mathcal{O}(\boldsymbol{\delta}^3) = \sqrt{\varepsilon_{1i}^2 + \varepsilon_{2i}^2}\ . \tag{H.5}$$

Next, inserting $(*1)$ into (H.3), we extract the non-oscillating terms at the inf. order $\boldsymbol{\delta}^2$. The result is

$$E = \frac{1}{2}m\mathbf{v}^2 + \frac{1}{4}m\omega^2\mathbf{A}_1^2 - \frac{c_{2b}^{3D}\hbar^2\omega^4\mathbf{A}^2}{2m\mathbf{v}^4} + \mathcal{O}(\boldsymbol{\delta}^3) = \frac{1}{2}m\mathbf{v}^2 + \frac{1}{2}m\omega^2\mathbf{A}_1^2 + \mathcal{O}(\boldsymbol{\delta}^3)\ . \tag{H.6}$$

From (H.5) and (H.6), we have

$$v_i = \frac{\sqrt{2mE_i}}{m}\big(\sqrt{1 + \varepsilon_{1i}^2 + \varepsilon_{2i}^2} - \sqrt{\varepsilon_{1i}^2 + \varepsilon_{2i}^2}\,\big) + \mathcal{O}(\boldsymbol{\delta}^3) = \frac{\sqrt{2mE_i}}{m}\big(1 - \sqrt{\varepsilon_{1i}^2 + \varepsilon_{2i}^2}\,\big) + \mathcal{O}(\boldsymbol{\delta}^3)\ . \tag{H.7}$$

Inserting (H.7) into (H.4), we have

$$c_{2b}^{3D} = -\frac{1}{2}\Big(1 - \frac{2\sum_{i=1}^{3} E_i\sqrt{\varepsilon_{1i}^2 + \varepsilon_{2i}^2}}{E}\Big) + \mathcal{O}(\boldsymbol{\varepsilon}^{\frac{3}{2}})\ .$$

Appendix I. Explicit Forms of $\mathsf{M}_1, \mathsf{N}_1, \mathsf{M}_2$, and N_2

We rewrite (3.8) to an easy-to-handle form. We expand $\tilde{s}$ in (3.8) with a power series of $\tilde{\xi}$ and $\tilde{\eta}$, then, takes the terms up to the second order (Fresnel approximation)[y] to have

$$\tilde{s} \simeq \tilde{s}\,' - (\tilde{l}\tilde{\xi} + \tilde{m}\tilde{\eta}) + \frac{(1-\tilde{l}\,^2)\tilde{\xi}^2 + (1-\tilde{m}^2)\tilde{\eta}^2 - 2\tilde{l}\tilde{m}\tilde{\xi}\tilde{\eta}}{2\tilde{s}\,'} \ ,$$

where $\tilde{l} = \tilde{X}/\tilde{s}\,',\ \tilde{m} = \tilde{Y}/\tilde{s}\,'$. We consider only the case $\tilde{m} = \tilde{Y}/\tilde{s}\,' = 0$. In this case, (3.8) becomes

$$\begin{aligned}
\tilde{U}(P) &= -\frac{iA}{\lambda\tilde{s}\,'}\int_{-\infty}^{-\mathsf{d}/2} d\tilde{\xi}\int_{-\infty}^{+\infty} d\tilde{\eta}\,\exp(-\frac{(\tilde{\xi}-\tilde{x}_0)^2+(\tilde{\eta}-\tilde{y}_0)^2}{w_0^2} + ik\tilde{s}) \\
&= -\frac{iA}{\lambda\tilde{s}\,'}\exp ik\tilde{s}\,'\int_{-\infty}^{-\mathsf{d}/2} d\tilde{\xi}\int_{-\infty}^{+\infty} d\tilde{\eta}\,\exp(-\frac{(\tilde{\xi}-\tilde{x}_0)^2}{w_0^2} + ik(-\tilde{l}\tilde{\xi} + \frac{(1-\tilde{l}\,^2)\tilde{\xi}^2}{2\tilde{s}\,'})) \\
&\qquad\qquad\qquad\qquad \times \exp(-\frac{(\tilde{\eta}-\tilde{y}_0)^2}{w_0^2} + ik\frac{\tilde{\eta}^2}{2\tilde{s}\,'})\ . \qquad \text{(I.1)}
\end{aligned}$$

We transform (I.1) to a tractable form. Setting $\tilde{\xi} = \tilde{\zeta}/\sqrt{\tilde{\alpha}} + k\tilde{l}/(\pi\tilde{\alpha})$, $\tilde{\alpha} = k(1-\tilde{l}^2)/(\pi\tilde{s}\,')$, we rewrite the real and imaginary parts of the $\tilde{\xi}$-direction integral as

$$\begin{aligned}
&\text{Re part:} \quad \int_{-\infty}^{-\mathsf{d}/2} d\tilde{\xi}\,\exp(-\frac{(\tilde{\xi}-\tilde{x}_0)^2}{w_0^2})\cos k(-\tilde{l}\tilde{\xi} + \frac{(1-\tilde{l}^2)\tilde{\xi}^2}{2\tilde{s}\,'}) \\
&= \frac{1}{\sqrt{\tilde{\alpha}}}\int_{-\infty}^{-\frac{\mathsf{d}\sqrt{\tilde{\alpha}}}{2}-\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}} d\tilde{\zeta}\,\exp(-\frac{(\tilde{\zeta}+\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}-\tilde{x}_0\sqrt{\alpha})^2}{\tilde{\alpha}w_0^2})\cos(\frac{\pi\tilde{\zeta}^2}{2} - \frac{k^2\tilde{l}\,^2}{2\pi\tilde{\alpha}}) \\
&= \cos(\frac{k^2\tilde{l}\,^2}{2\pi\tilde{\alpha}})\cdot\frac{1}{\sqrt{\tilde{\alpha}}}\int_{-\infty}^{-\frac{\mathsf{d}\sqrt{\tilde{\alpha}}}{2}-\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}} d\tilde{\zeta}\,\exp(-\frac{(\tilde{\zeta}+\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}-\tilde{x}_0\sqrt{\tilde{\alpha}})^2}{\tilde{\alpha}w_0^2})\cos\frac{\pi\tilde{\zeta}^2}{2} \\
&+ \sin(\frac{k^2\tilde{l}\,^2}{2\pi\tilde{\alpha}})\cdot\frac{1}{\sqrt{\tilde{\alpha}}}\int_{-\infty}^{-\frac{\mathsf{d}\sqrt{\tilde{\alpha}}}{2}-\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}} d\tilde{\zeta}\,\exp(-\frac{(\tilde{\zeta}+\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}-\tilde{x}_0\sqrt{\tilde{\alpha}})^2}{\tilde{\alpha}w_0^2})\sin\frac{\pi\tilde{\zeta}^2}{2} \\
&= \cos(\frac{k^2\tilde{l}\,^2}{2\pi\tilde{\alpha}})\cdot\frac{1}{\sqrt{\tilde{\alpha}}}\mathsf{M}_1 + \sin(\frac{k^2\tilde{l}\,^2}{2\pi\tilde{\alpha}})\cdot\frac{1}{\sqrt{\tilde{\alpha}}}\mathsf{N}_1\ ,
\end{aligned}$$

---

[y] Because (tildes omitted)

$$\sqrt{Z^2+(X-\xi)^2} = s' - \xi l + \frac{\xi^2(1-l^2)}{2s'} + \frac{\xi^3 l(1-l^2)}{2s'^2} + \frac{\xi^4}{s'^3}(-\frac{1}{8}+\frac{3l^2}{4}-\frac{5l^4}{8}) + \cdots\ ,$$

if

$$k|\frac{\xi^3 l(1-l^2)}{2s'^2} + \frac{\xi^4}{s'^3}(-\frac{1}{8}+\frac{3l^2}{4}-\frac{5l^4}{8}) + \cdots| << 2\pi \quad \cdots(*)$$

is satisfied at P, the Fresnel approximation is effective at the P (e.g., Ref. 21 §8.3.3). In the present case, setting the criteria as $k|\cdots| < 2\pi \times 0.05$, the Fresnel approximation is effective in $7.3mm < Z$ if $X < 0.3Z$, effective in $4.5mm < Z$ if $X < 0.1Z$, effective in $0.6mm < Z$ if $X = 0$. In each case, the moving range of ξ was set to $-\sqrt{2}w_0 < \xi < \sqrt{2}w_0$.

$$\text{Im part:}\quad \int_{-\infty}^{-\mathsf{d}/2} d\tilde{\xi}\exp\Big(-\frac{(\tilde{\xi}-\tilde{x}_0)^2}{w_0^2}\Big)\sin k\Big(-\tilde{l}\tilde{\xi}+\frac{(1-\tilde{l}^2)\tilde{\xi}^2}{2\tilde{s}\,'}\Big)$$
$$=\frac{1}{\sqrt{\tilde{\alpha}}}\int_{-\infty}^{-\frac{\mathsf{d}\sqrt{\tilde{\alpha}}}{2}-\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}} d\tilde{\zeta}\exp\Big(-\frac{(\tilde{\zeta}+\frac{k\tilde{l}}{2\sqrt{\tilde{\alpha}}}-\tilde{x}_0\sqrt{\tilde{\alpha}})^2}{\tilde{\alpha}w_0^2}\Big)\sin\Big(\frac{\pi\tilde{\zeta}^2}{2}-\frac{k^2\tilde{l}^{\,2}}{2\pi\tilde{\alpha}}\Big)$$
$$=\cos\Big(\frac{k^2\tilde{l}^{\,2}}{2\pi\tilde{\alpha}}\Big)\cdot\frac{1}{\sqrt{\tilde{\alpha}}}\mathsf{N}_1-\sin\Big(\frac{k^2\tilde{l}^{\,2}}{2\pi\tilde{\alpha}}\Big)\cdot\frac{1}{\sqrt{\tilde{\alpha}}}\mathsf{M}_1\ .$$

where M_1 and N_1 are

$$\mathsf{M}_1=\int_{-\infty}^{-\frac{\mathsf{d}\sqrt{\tilde{\alpha}}}{2}-\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}} d\tilde{\zeta}\exp\Big(-\frac{(\tilde{\zeta}+\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}-\tilde{x}_0\sqrt{\tilde{\alpha}})^2}{\tilde{\alpha}w_0^2}\Big)\cos\frac{\pi\tilde{\zeta}^2}{2}$$
$$\simeq\int_{-\infty}^{-\sqrt{\frac{k}{\pi\tilde{Z}}}(\frac{\mathsf{d}}{2}+\tilde{X})} d\tilde{\zeta}\exp\Big(-\frac{(\tilde{\zeta}+\sqrt{\frac{k}{\pi\tilde{Z}}}(\tilde{X}-\tilde{x}_0))^2}{\frac{kw_0^2}{\pi\tilde{Z}}}\Big)\cos\frac{\pi\tilde{\zeta}^2}{2}\ ,$$

$$\mathsf{N}_1=\int_{-\infty}^{-\frac{\mathsf{d}\sqrt{\tilde{\alpha}}}{2}-\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}} d\tilde{\zeta}\exp\Big(-\frac{(\tilde{\zeta}+\frac{k\tilde{l}}{\pi\sqrt{\tilde{\alpha}}}-\tilde{x}_0\sqrt{\tilde{\alpha}})^2}{\tilde{\alpha}w_0^2}\Big)\sin\frac{\pi\tilde{\zeta}^2}{2}$$
$$\simeq\int_{-\infty}^{-\sqrt{\frac{k}{\pi\tilde{Z}}}(\frac{\mathsf{d}}{2}+\tilde{X})} d\tilde{\zeta}\exp\Big(-\frac{(\tilde{\zeta}+\sqrt{\frac{k}{\pi\tilde{Z}}}(\tilde{X}-\tilde{x}_0))^2}{\frac{kw_0^2}{\pi\tilde{Z}}}\Big)\sin\frac{\pi\tilde{\zeta}^2}{2}\ .$$

Likewise, setting $\tilde{\eta}=\tilde{\rho}\sqrt{\pi\tilde{s}\,'/k}$, we write the real and imaginary parts of the $\tilde{\eta}$-direction integral as

$$\text{Re part:}\quad \int_{-\infty}^{+\infty} d\tilde{\eta}\exp\Big(-\frac{(\tilde{\eta}-\tilde{y}_0)^2}{w_0^2}\Big)\cos\frac{k\tilde{\eta}^2}{2\tilde{s}\,'}$$
$$=\sqrt{\frac{\pi\tilde{s}\,'}{k}}\int_{-\infty}^{+\infty} d\tilde{\rho}\exp\Big(-\frac{(\tilde{\rho}-\frac{\tilde{y}_0}{\sqrt{\pi\tilde{s}\,'/k}})^2}{\frac{kw_0^2}{\pi\tilde{s}\,'}}\Big)\cos\frac{\pi\tilde{\rho}^2}{2}=\sqrt{\frac{\pi\tilde{s}\,'}{k}}\ \mathsf{Q}\ ,$$

$$\text{Im part:}\quad \int_{-\infty}^{+\infty} d\tilde{\eta}\exp\Big(-\frac{(\tilde{\eta}-\tilde{y}_0)^2}{w_0^2}\Big)\sin\frac{k\tilde{\eta}^2}{2\tilde{s}\,'}$$
$$=\sqrt{\frac{\pi\tilde{s}\,'}{k}}\int_{-\infty}^{+\infty} d\tilde{\rho}\exp\Big(-\frac{(\tilde{\rho}-\frac{\tilde{y}_0}{\sqrt{\pi\tilde{s}\,'/k}})^2}{\frac{kw_0^2}{\pi\tilde{s}\,'}}\Big)\sin\frac{\pi\tilde{\rho}^2}{2}=\sqrt{\frac{\pi\tilde{s}\,'}{k}}\ \mathsf{R}\ ,$$

in which Q and R represent the integrals of the left sides. For the present purpose, we can set $y_0=0$. Then, because $kw_0^2/\pi\tilde{s}\,'$ is large in the present case, the approximate values of Q and R are obtained from the Fresnel integrals: $\int_0^\infty\cos(\pi\tau^2/2)d\tau=1/2,\int_0^\infty\sin(\pi\tau^2/2)d\tau=$

$1/2$. That is, $\mathsf{Q} \simeq 1, \mathsf{R} \simeq 1$. Thus, (I.1) is written as (note $-\frac{iA}{\lambda \tilde{s}\,'} \frac{1}{\sqrt{\tilde{\alpha}}} \sqrt{\frac{\pi \tilde{s}\,'}{k}}(1+i) = \frac{Ae^{-i\pi/4}}{\sqrt{2(1-\tilde{l}^2)}}$)

$$\tilde{U}(P) = \frac{Ae^{-i\pi/4}}{\sqrt{2(1-\tilde{l}^2)}} e^{ik\tilde{s}\,'}$$
$$\times \Big((\cos(\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}}) \cdot \mathsf{M}_1 + \sin(\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}}) \cdot \mathsf{N}_1) + i(\cos(\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}}) \cdot \mathsf{N}_1 - \sin(\frac{k^2 \tilde{l}^{\,2}}{2\pi\tilde{\alpha}}) \cdot \mathsf{M}_1) \Big) \ .$$

In a same manner, $\tilde{\tilde{U}}(P)$ is obtained as

$$\tilde{\tilde{U}}(P) = \frac{Ae^{-i\pi/4}}{\sqrt{2(1-\tilde{\tilde{l}}^2)}} e^{ik\tilde{\tilde{s}}\,'}$$
$$\times \Big((\cos(\frac{k^2 \tilde{\tilde{l}}^{\,2}}{2\pi\tilde{\tilde{\alpha}}}) \cdot \mathsf{M}_2 + \sin(\frac{k^2 \tilde{\tilde{l}}^{\,2}}{2\pi\tilde{\tilde{\alpha}}}) \cdot \mathsf{N}_2) + i(\cos(\frac{k^2 \tilde{\tilde{l}}^{\,2}}{2\pi\tilde{\tilde{\alpha}}}) \cdot \mathsf{N}_2 - \sin(\frac{k^2 \tilde{\tilde{l}}^{\,2}}{2\pi\tilde{\tilde{\alpha}}}) \cdot \mathsf{M}_2) \Big) \ ,$$

where

$$\mathsf{M}_2 = \int_{\frac{d\sqrt{\tilde{\tilde{\alpha}}}}{2} - \frac{k\tilde{\tilde{l}}}{\pi\sqrt{\tilde{\tilde{\alpha}}}}}^{+\infty} d\tilde{\tilde{\zeta}} \exp(-\frac{(\tilde{\tilde{\zeta}} + \frac{k\tilde{\tilde{l}}}{\pi\sqrt{\tilde{\tilde{\alpha}}}} - \tilde{\tilde{x}}_0\sqrt{\tilde{\tilde{\alpha}}})^2}{\tilde{\tilde{\alpha}} w_0^2}) \cos\frac{\pi\tilde{\tilde{\zeta}}^2}{2}$$
$$\simeq \int_{\sqrt{\frac{k}{\pi\tilde{\tilde{Z}}}}(\frac{\mathsf{d}}{2} - \tilde{\tilde{X}})}^{+\infty} d\tilde{\tilde{\zeta}} \exp(-\frac{(\tilde{\tilde{\zeta}} + \sqrt{\frac{k}{\pi\tilde{\tilde{Z}}}}(\tilde{\tilde{X}} - \tilde{\tilde{x}}_0))^2}{\frac{kw_0^2}{\pi\tilde{\tilde{Z}}}}) \cos\frac{\pi\tilde{\tilde{\zeta}}^2}{2} \ ,$$
$$\mathsf{N}_2 = \int_{\frac{d\sqrt{\tilde{\tilde{\alpha}}}}{2} - \frac{k\tilde{\tilde{l}}}{\pi\sqrt{\tilde{\tilde{\alpha}}}}}^{+\infty} d\tilde{\tilde{\zeta}} \exp(-\frac{(\tilde{\tilde{\zeta}} + \frac{k\tilde{\tilde{l}}}{\pi\sqrt{\tilde{\tilde{\alpha}}}} - \tilde{\tilde{x}}_0\sqrt{\tilde{\tilde{\alpha}}})^2}{\tilde{\tilde{\alpha}} w_0^2}) \sin\frac{\pi\tilde{\tilde{\zeta}}^2}{2}$$
$$\simeq \int_{\sqrt{\frac{k}{\pi\tilde{\tilde{Z}}}}(\frac{\mathsf{d}}{2} - \tilde{\tilde{X}})}^{+\infty} d\tilde{\tilde{\zeta}} \exp(-\frac{(\tilde{\tilde{\zeta}} + \sqrt{\frac{k}{\pi\tilde{\tilde{Z}}}}(\tilde{\tilde{X}} - \tilde{\tilde{x}}_0))^2}{\frac{kw_0^2}{\pi\tilde{\tilde{Z}}}}) \sin\frac{\pi\tilde{\tilde{\zeta}}^2}{2} \ .$$

$\mathsf{M}_1, \mathsf{N}_1, \mathsf{M}_2$, and N_2 at $P = (X, 0, Z)$ on the coordinate system (x, y, z) are expressed explicitly, ignoring terms of the second and higher powers of X/Z, as

$$\mathsf{M}_1|_{P=(X,0,Z)} = \int_{-\infty}^{\sqrt{\frac{k}{\pi Z}}(-\frac{\mathsf{d}}{2} - X + \frac{k_x}{k_z}Z)} \exp\Big(- \frac{\big(\tau + \sqrt{\frac{k}{\pi Z}}(X - \frac{k_x}{k_z}Z - x_0)\big)^2}{\frac{kw_0^2}{\pi Z}} \Big) \cos\frac{\pi\tau^2}{2} d\tau \tag{I.2a}$$

$$\mathsf{N}_1|_{P=(X,0,Z)} = \int_{-\infty}^{\sqrt{\frac{k}{\pi Z}}(-\frac{\mathsf{d}}{2} - X + \frac{k_x}{k_z}Z)} \exp\Big(- \frac{\big(\tau + \sqrt{\frac{k}{\pi Z}}(X - \frac{k_x}{k_z}Z - x_0)\big)^2}{\frac{kw_0^2}{\pi Z}} \Big) \sin\frac{\pi\tau^2}{2} d\tau \ . \tag{I.2b}$$

$$\mathsf{M}_2|_{P=(X,0,Z)} = \int_{\sqrt{\frac{k}{\pi Z}}(+\frac{\mathsf{d}}{2}-X-\frac{k_x}{k_z}Z)}^{+\infty} \exp\Big(- \frac{\big(\tau + \sqrt{\frac{k}{\pi Z}}(X + \frac{k_x}{k_z}Z - x_0)\big)^2}{\frac{k w_0^2}{\pi Z}}\Big) \cos\frac{\pi\tau^2}{2} d\tau \quad \text{(I.3a)}$$

$$\mathsf{N}_2|_{P=(X,0,Z)} = \int_{\sqrt{\frac{k}{\pi Z}}(+\frac{\mathsf{d}}{2}-X-\frac{k_x}{k_z}Z)}^{+\infty} \exp\Big(- \frac{\big(\tau + \sqrt{\frac{k}{\pi Z}}(X + \frac{k_x}{k_z}Z - x_0)\big)^2}{\frac{k w_0^2}{\pi Z}}\Big) \sin\frac{\pi\tau^2}{2} d\tau\ . \quad \text{(I.3b)}$$

References

1. W. Greiner, *Quantum mechanics: an introduction*, 4th edn. (Springer-Verlag, 2000).
2. P. R. Holland, *The Quantum Theory of Motion* (Cambridge University Press, 1993).
3. A. E. Faraggi and M. Matone, *Int. J. Mod. Phys. A* **15** (13), 1869-2017 (2000).
4. A. Bouda, *Int. J. Mod. Phys. A* **18** (19), 3347-3368 (2003).
5. R. Miron, *The Geometry of Higher-Order Lagrange Spaces* (Kluwer A. P., 1997).
6. M. de León and P. R. Rodrigues, *Generalized Classical Mechanics and Field Theory* (North-Holland, 1985).
7. J. Komrska, V. Drahoš and A. Delong, *Optica Acta* **14** (2), 147-167 (1967).
8. A. Ohshita, H. Teraoka, Y.Mametani and H. Tomita, *J. Electron Microsc.* **35** (2), 111-117 (1986).
9. K. Yamamoto, T. Tanji and M. Hibino, *Ultramicroscopy* **85**, 35-49 (2000).
10. V. Ovsienko and S. Tabachnikov, *Projective differential geometry old and new: from the Schwarzian derivative to the cohomology of diffeomorphism groups* (Cambridge U. P., 2005).
11. K. S. Brown, *Cohomology of Groups* (Springer-Verlag New York, Inc., 1982).
12. D. J. S. Robinson, *A Course in the Theory of Groups*, 2nd edn. (Springer-Verlag, 1996).
13. J. J. Rotman, *An Introduction to the Theory of Groups*, 4th edn. (Springer-Verlag, 1995).
14. A. Messiah, *Quantum mechanics* (North Holland Publishing Company Amsterdam, 1961-1962).
15. I. Anderson and G. Thompson, *The Inverse Problem of the Calculus of Variations for Ordinary Differential Equations* (Mem. Am. Math. Soc. **98** (473), 1992).
16. E. A. Coddington, *An Introduction to Ordinary Differential Equations* (Dover, 1989).
17. N. N. Lebedev, *Special Functions & Their Applications* (Dover, 1972).
18. J. Durnin, J. J. Miceli and J. H. Eberly, *Phys. Rev. Lett.* **58** (15), 1499-1501 (1987).
19. O. Svelto, *Principles of Lasers*, 4th edn. (Plenum Press, New York, 1998).
20. J. E. Pearson, T. C. McGill, S. Kurtin and A. Yariv, *J. Opt. Soc. Am.* **59** (11), 1440-1445 (1969).
21. M. Born and E. Wolf, *Principles of Optics*, 7th edn. (Cambridge University Press, 1999).
22. L. D. Landau and E. M. Lifshitz, *The classical theory of fields*, 4th rev. English edn. (Pergamon Press, 1975).
23. G. Bertoldi, A. E. Faraggi and M. Matone, *Class.Quantum Grav.* **17**, 3965-4005 (2000).
24. M. de León and P. R. Rodrigues, *Methods of Differential Geometry in Analytical Mechanics* (North-Holland, 1989).
25. M. Crampin, W Sarlet and F. Cantrijn, Math. Proc. Camb. Phil. Soc. **99** 565-587 (1986).